

PROPOSAL FORM

THIS FORM MUST BE SUBMITTED BY THE PROPOSED PRINCIPAL INVESTIGATOR (S) AND SUBMITTED ALONG WITH THE PROPOSAL.

By submission of this proposal, I agree to abide by the Trustee Council's data policy (*Trustee Council Data Policy**, adopted March 17, 2008) and reporting requirements (*Procedures for the Preparation and Distribution of Reports***, adopted June 27, 2007).

PROJECT TITLE: Amendment to Lingering Oil on Boulder-Armored Beaches in the Gulf of Alaska 23 Years after the Exxon Valdez Oil Spill (Project 11100112)

Name of PI: Gail V. Irvine
Email: girvine@usgs.gov Phone: 907-786-7069
Mailing Address U.S. Geological Survey, 4210 University Drive
City, State, Zip Anchorage, AK 99508

Name of PI: Daniel H. Mann
Email: dhmann@alaska.edu Phone: 907-474-7127
Mailing Address Scenarios Network for Alaska Planning, Geography Program
University of Alaska
City, State, Zip Fairbanks, AK 99775

Name of PI: Mark G. Carls
Email: mark.carls@noaa.gov Phone: 907-789-6019
Mailing Address National Marine Fisheries Service, Auke Bay Laboratories
17109 Pt. Lena Loop Road
City, State, Zip Juneau, AK 99801

* www.evostc.state.ak.us/Policies/data.cfm

** www.evostc.state.ak.us/Policies/reporting.cfm

FY13 AMENDMENT PROPOSAL SUMMARY PAGE

Project Title: Amendment to Lingering Oil on Boulder-Armored Beaches in the Gulf of Alaska 23 Years after the *Exxon Valdez* Oil Spill (Project 11100112)

Project Period: FFY2011-2012-2013

Primary Investigator(s): Dr. Gail Irvine (USGS), Dr. Daniel Mann (Hovawart Mann Works), Mark Carls (NOAA, NMFS)

Study Location: Gulf of Alaska, (Katmai National Park & Preserve, Kenai Fjords NP&P)

Amendment Abstract: This amendment to Project 11100112 solely requests funding for additional hydrocarbon analyses. Our request is prompted by: 1) findings from chemical analyses of passive samplers that were deployed in 2011, and 2) the opportunity to obtain additional samples in 2012 because the majority of our field work was delayed by weather until this summer. The findings from 2011 indicated the presence of dissolved oil constituents at one of the two oiled sites where we deployed passive samplers. There was also some visual indication that the boulder armor at that site had been disturbed. Redeploying passive samplers at the same oiled and control sites this year, prior to our more extensive sampling via chartered vessel, will allow us to make within-site and between-site assessments of oil release over two years. We have sufficient funds for the logistic costs, therefore this is a potentially very high-benefit, low-cost proposition. In this amendment, we are requesting funds for the analysis of hydrocarbon samples by comprehensive two-dimensional gas chromatography (GCxGC); these would be analyzed by Woods Hole Oceanographic Institution (Dr. Chris Reddy). This analytical technique offers an order of magnitude improvement in detection limit and has greater ability to discriminate components of the remaining hydrocarbons than traditional gas chromatography (GC). GCxGC-based techniques use all GC amenable compounds within an oil to examine weathering trends, not just those compounds whose identities are known (or visible using one-dimensional chromatography). We feel it would make a very worthwhile addition to this project. There are no changes to objectives nor to due dates under this amendment. The accompanying budget details our funding request.

Original Project Abstract: We want to continue long-term monitoring of lingering oil at six Gulf of Alaska sites where we have tracked the fate and persistence of stranded *Exxon Valdez* oil over the last 22 years. It has been six years since our last survey revealed that relatively unweathered oil still persisted at some sites. Interestingly these sites have less weathered oil (e.g., contains more *n*-alkanes) than similarly aged oil from Prince William Sound. All five of our monitoring sites on the Katmai National Park coast are boulder beaches with high wave energies. Accepted knowledge predicted that rapid natural weathering of stranded oil would occur in such settings. This was not the case, and we are still figuring out why. We think it is because the boulder armors that cover these shorelines protect the underlying oil. In addition to resampling our monitoring plots, we will be testing to see if oil is leaking out from these beaches. By extending our long term study of oil stranded on this little understood shoreline type, we will contribute important new data useful for predicting the geographic distribution of lingering oil, assessing its potential for continued pollution, and designing methods for its remediation.

Estimated Budget: New: \$28.4k + 9%GA = \$31.0k

EVOS Funding Requested: FFY2013 (New): \$31.0k

Non-EVOS Funds to be used: FFY2013: \$26.0k

Date: June, 2012

BUDGET JUSTIFICATION – Note: Values discussed here reflect only the additional funds needed and requested under this amendment

Personnel: New amount requested FY13 - \$ 0k

Travel: New amount requested FY13– \$ 0k

Contractual: New amount requested FY13- \$ 15.6k

All contractual costs are for GCxGC analysis of hydrocarbons from sediment samples by Woods Hole Oceanographic Institution, Dr. Chris Reddy: 13 samples @ \$1200/sample = \$15,600

Commodities: New amount requested FY13- \$ 12.8k

The commodities cost, \$12.8k, is for NOAA's analysis of the LDPEs deployed and retrieved this year (FY2012) under this amendment. Construction of passive samplers (LDPEs), and GC-MS analysis of the plastic strips for hydrocarbons (PAHs): 30 samplers @ \$425 = \$12,750 or \$12.8K

Equipment: New amount requested FY13- \$ 0k

TOTAL REQUESTED: \$28,400 plus \$2,600 GA = \$31,000

13100112 Amendment: Agency Distribution

USGS: \$15,600 plus \$1400 GA = \$17,000

NOAA: \$12,800 plus \$1200 GA = \$14,000

Non-EVOS Funds to be used: FFY2013: \$26.0k

In-kind contributions:

NOAA - hydrocarbon analytical support: \$20K

Mark Carls, 1 month @ \$8k/mo= \$8k

Jeep Rice, 0.5 month @ \$10k/mo= \$5k

Mandy Lindeberg, 1 month @ \$7k/mo=\$7k

USGS: Gail Irvine, 0.5 mo @ \$11.7= \$6k

Lingering Oil on Boulder Armored Beaches in the Gulf of Alaska 23 Years after the *Exxon Valdez* Oil Spill

PROJECT PLAN

I. NEED FOR THE PROJECT

A. Statement of Problem

Contrary to the predictions made by oil-spill experts in 1989, significant amounts of *Exxon Valdez* oil remain in the spill region 20 years later. Short et al. (2004) estimate there are 7.8 hectares of oiled shorelines left in Prince William Sound (PWS) containing some 56,000 kilograms (kg) of lingering oil in the subsurface. Furthermore, Short et al. (2007) assert that the areal extent of oiled beaches in PWS did not change significantly between 2001 and 2005, which implies that the rate of decline in lingering oil has slowed. There have been no detailed studies of the amount of lingering oil in the Gulf of Alaska; however, long-term monitoring of oiled sites shows persistence of relatively unweathered oil in Kenai Fjords National Park and Katmai National Park (Irvine et al., 1999; 2006; 2007). Is that oil still there on high wave energy beaches in the Gulf of Alaska? Has its chemical weathering changed significantly? Is it leaking from the beach, thereby potentially posing biological threat? If it is still there, what factors are causing it to persist?

1) Background

a) Lingering *Exxon Valdez* Oil

Our knowledge about lingering oil in the spill region has become much more complete over the last 20 years, but large gaps still exist. Some of these gaps involve geographical differences in oiling and geomorphology/exposure within the spill region. For example, in PWS oil reached shorelines in a more fluid or less viscous form than the emulsified water/oil form (mousse) that landed on GOA shores. Also, PWS is in general a more protected environment than the GOA, and this basic fact has widespread implications for the coastal geomorphologies of the two areas. We know now that in PWS much of the remaining oil is found at a lower level in the intertidal zone than was thought immediately after the spill (Short et al., 2006). On the other hand, this is not a universal pattern, since lingering oil in the GOA tends to be located high in the intertidal zone (Irvine et al., 2006). A modeling study designed to predict the location of lingering subsurface oil within the spill area, including both Prince William Sound (PWS) and the Gulf of Alaska (GOA), was begun in 2007 (Michel et al., 2010). This ground-breaking work has developed geospatial models that identify areas where subsurface oil is still present on the shorelines of PWS and the GOA and estimate the relative quantities of subsurface oil remaining at different sites. One of the most significant results of this work is its prediction that a significant number of as-yet-unsurveyed sites in PWS and the GOA still contain subsurface oil. On the down side, this geospatial model has been developed primarily based on data from PWS and so has limited applicability to GOA sites. Furthermore, it is implicit in multi-variable models that while overall predictive success may be high (as in PWS), the linkages between the data used and the physical phenomena that drive oil persistence remain unclear (Michel et al., 2010). In other words, the model may work, but we still do not understand the geomorphic and geochemical processes that allow the persistence of stranded oil.

Other recent EVOS-funded studies focus on smaller-scale processes related to subsurface oil persistence. M. Boufadel and collaborators are studying factors that limit the degradation rate of oil in PWS beaches including nutrient and oxygen concentrations and water flow (Boufadel et al., 2010; Li and Boufadel, 2010). A. Venosa et al. (2010) have researched the factors limiting biodegradability of oiled sediment. Both these small-scale, process studies emphasize the importance of oxygenation, nutrient availability, and hydraulic conductivity in the subsurface of oiled beaches. Certainly, these small-scale variables are influenced at larger spatial scales by the nature and stability of the overlying armor layers.

b) Boulder Armored Beaches

Boulder armors develop naturally when the finer particles (silt, sand, pebbles, and cobbles) are winnowed away by waves, deflating the pre-existing sediments until a layer of boulders remains that prevents further winnowing. Natural boulder armors are little studied despite their wide distribution on shorelines around the world and despite the widespread use of artificial boulder armors to stabilize eroding beaches (Dean and Dalrymple, 2004). Natural and artificial armors are distinctly different phenomena, and the stability formulae used to design artificial armors have little relevance to natural armors (Oak, 1986). A recent review of armored, gravel beaches on paraglacial coastlines is given by Hayes et al. (2010).

Boulder beaches are often intricately packed or fitted together with the projections of one boulder accommodated in the concavities of its neighbors (Shelley, 1968). Smaller boulders are often imprisoned amongst larger ones (Hills, 1970). The fitting together of boulder armors occurs by boulders shifting in place, rubbing against their neighbors until achieving a packing of maximum stability. Tracking of the positions of individually marked boulders on the Katmai coastline shows that while individual boulders regularly roll and shift in place, few ever move out of their niches within the surrounding armor (Irvine et al., 2006; 2007). Armors form tightly fitted fabrics that are highly resistant to wave attack and may be stable for thousands of years (Bishop and Hughes, 1989). Hence boulder armors represent equilibrium geomorphic features; that is, they develop into progressively more stable entities to the point where most wave events cannot disturb them or the sediments (and oil) they cover. Boulder armors are ubiquitous on Gulf of Alaska shorelines (Hayes et al., 2010). Exceptions are shorelines where sea-level changed radically during the Great Alaskan Earthquake in 1964 and shorelines experiencing rapid progradation by glacial outwash.

In summary, naturally occurring boulder armors are widespread on rocky shorelines. Because they are created through waves, armors are most common and best developed on high energy shorelines like many in the GOA and on exposed shorelines in PWS. The dynamics of boulder armors have been little studied relative to sandy and gravel beaches, which tend to be more widespread at lower latitudes. As a result, the processes important in the development and maintenance of boulder armors remain poorly known, though it is clear that boulder beaches are quite different from sand and gravel beaches with a unique set of formative processes (Oak, 1984; Hayes et al., 2010). Another thing that is clear is that boulder-armored shorelines can harbor slightly weathered oil for long periods of time (Irvine et al., 2006; 2007; Short et al., 2007). It seems likely that if there is still *Exxon Valdez* oil in the environment of southern Alaska 50 years hence, it will be associated with boulder armors.

c) Our Long Term Monitoring Study of GOA Shorelines

Since 1994, we have monitored the status of *Exxon Valdez* oil at six sites in the Gulf of Alaska (Irvine et al., 1999; 2006; 2007; Short et al., 2007). These sites are now the most consistently studied, long-term monitoring sites of stranded oil in the spill region. Sixteen years post-spill, surface oiling had declined markedly at all sites, but subsurface oil remained abundant. The oil collected from beneath the boulder armor at three of the four sites surveyed was still compositionally similar to eleven-day old *Exxon Valdez* oil (Short et al., 2007). Remarkably, this oil still contained *n*-alkanes, which normally would be degraded by microbes within weeks of a spill. When the composition of *Exxon Valdez* oil from the GOA was compared to that from PWS, the GOA oil was less-weathered (Short et al., 2007). These findings indicate that our GOA study of the long-term persistence of stranded oil may provide insights not possible from PWS studies and that may apply to some of the extensive coastline that was oiled outside of PWS.

The persistence of oil at high wave-energy sites in the GOA seems to be related to the presence of stable boulder armors. Though not initially chosen for this reason, all five of our monitoring sites on the Katmai National Park and Preserve coast in the GOA possess such boulder armors. The prediction that oil persistence correlates with armor stability has been borne out over the last 16 years. Analysis of movements in the boulder armors reveals that only minor shifts have occurred since 1994. These findings suggest that boulder armors, combined with the stranding of oil mousse high in the intertidal zone, results in the unexpectedly lengthy persistence of only slightly to moderately weathered oil within otherwise high-energy wave environments on GOA coastlines. The three-dimensional matrix provided by boulder-armored beaches allows oil to penetrate into finer sediments lying beneath stable, boulder lags. Previously it was thought that oil would be rapidly removed from such geomorphic settings by the vigorous wave action (Vandermuelen, 1977). Instead, these surface armors attenuate wave energy and reduce wave reworking of the underlying substrates and the included oil. Additionally, oil on boulder-armored beaches is sheltered by the boulders from sun exposure (Irvine et al., 1999). Similar inferences about the importance of boulder armors in allowing oil to persist for long periods on exposed shorelines comes from observations made inside PWS (Michel and Hayes, 1993a, b; 1995; 1999; Hayes and Michel, 1999; Hayes et al., 2010). Understanding the dynamics of armored shorelines is basic to understanding what determines the distribution of persistent, subsurface oil.

The persistence of this oil in the GOA raises questions about its potential or realized biological effects. In PWS a number of studies have examined biological effects of the spill over the years (e.g., Bodkin et al., 2002; Esler and Iverson, 2010), but these types of studies are lacking in the GOA except for more limited temporal sampling of oiled mussel beds (Babcock, et al., 1996; Carls, et al., 2001, Irvine et al., 2007). Thus the ability to tie lingering oil to biotic effects is limited. We propose to examine whether oil is being released from these sites as a first step in addressing this particular gap in our understanding of biological effects of lingering oil.

B. Relevance to 1994 Restoration Plan Goals and Scientific Priorities

Our proposed work will address the physical and chemical processes responsible for the persistence of lingering oil in the spill region within the GOA and seeks to understand the reasons why this long lingering oil has failed to degrade. Additionally, we are investigating whether the oil is being released and may be affecting biota. Of particular significance is the fact that five of our long-term monitoring sites are located within a designated wilderness area in

Katmai National Park and Preserve. Our findings will provide direct evidence of the recovery status of these special-value lands and will assist in the evaluation of remediation options that could lead to restoration of these injured natural resources. Our proposed study of lingering subsurface oil on boulder armored beaches in the GOA will fill a geographical gap in our understanding of the distribution of lingering oil and directly complement recent or ongoing studies of oil biodegradation at finer spatial scales.

II. PROJECT DESIGN

Objective #1. What is the status of oiling at our long-term monitoring sites, 23 years after the *Exxon Valdez* spill? Specifically, how chemically weathered is the oil today, and how have the extents of surface and subsurface oiling changed?

Objective #2: How much of the subsurface oil preserved under boulder armors at our GOA monitoring sites is presently leaking into the surrounding environment?

Objective #3: How stable have the boulder armors on our study beaches been over the last 23 years and how does this relate to the findings from Objectives #1 and 2?

B. Procedural and Scientific Methods

1) What is the status of oiling at our long-term monitoring sites, 23 years after the *Exxon Valdez* spill? Specifically, how weathered is the oil and how have the extents of surface and subsurface oiling changed?

We will reassess the extent of both surface and subsurface oil using the same methods we have used since 1994 at these sites. Additionally, we will collect two oiled sediment samples from each site for hydrocarbon analyses. These samples will be analyzed via gas-chromatography/mass-spectrometry (GCMS) by NOAA's Auke Bay Laboratory.

2) Is the subsurface oil preserved under boulder armors presently leaking into surrounding environment?

Although oil has persisted at our GOA monitoring sites for at least 16 years, we do not know if oil is presently leaking from the subsurface into the environment. If it is occurring, such leakage could be having biological impacts. To ascertain if oil is leaking out, we will deploy low density polyethylene strips (LDPEs), which we refer to here simply as "plastic strips." These plastic strips function like the better known semi-permeable membrane devices (SPMDs) (Chapman, 2006), but are superior when the hydrocarbon signal is low (e.g., in relatively unpolluted environments), since they record less background 'noise' than do SPMDs (Jeep Rice, pers. comm.). Polynuclear aromatic hydrocarbons (PAHs) are adsorbed onto the plastic strips, but not alkanes or particulate oil. We will deploy the plastic strips, in their protective containers, in radiating patterns near boulder armors that still shelter remnant oil, and also at control sites. Our plan is to place the plastic strips at two of our long-term monitoring sites on the Katmai coast and at two un-oiled control sites relatively near these oiled sites. At each oiled site, we will deploy 10 plastic strips, while at each control site we plan to deploy 4 plastic strips. Trip and field blanks will be collected and analyzed for control purposes. At all sites, the plastic strips will be left in place for up to 30 days, then collected for analysis of hydrocarbons. We also plan to collect mussels (*Mytilus trossulus*) near these same sites - where they are present - and analyze

them for hydrocarbons as well, since they are better indicators of particulate hydrocarbons (Jeff Short, pers. comm.).

3) How stable are the boulder armors?

We will resurvey the locations of the marked boulders at each site, using the same methods as previously. The deviations from the previous locations will be calculated and used to determine if individual boulders have moved significantly over time. The degree of boulder movement on each beach will be used to interpret the data gathered in Objectives #1 and #2 on the extent, chemical composition of oiling and whether oil is being released into the environment. If boulder armors are responsible for the long term persistence of EVOS oil, we expect to see the most oil and the least weathered oil at sites whose armors have moved the least.

C. Data Analysis and Statistical Methods

Surface oiling at our GOA monitoring sites is reassessed in marked quadrats by estimating oil percent cover. Percent cover data for individual quadrats will be compared through time (1994, 1999, 2005, and 2012) via pair-wise tests. As for all tests discussed here, the data will be tested for normality and the appropriate parametric or non-parametric test chosen. Data from previous years (1994, 1999 and 2005) were compared in our latest report and manuscript via Wilcoxon signed-rank tests.

Subsurface oiling is assessed through the sampling of “dip stones” at each site. These are naturally occurring cobbles that extend from the sub-armor surface of the substrate downwards through the zone of subsurface oiling. Means and ranges of the depth of oiling for each site will be compared through time.

Hydrocarbon analyses:

Oil composition and weathering: As in our previous studies, chemical analysis of sediment, mussel and LDPE samples will be conducted via gas-chromatography/mass-spectrometry (Short et al., 1996a). We will compare the presence and relative abundance of polynuclear aromatic hydrocarbons (PAH) within samples, and compute a weathering index based on a first-order kinetic loss rate model of Short and Heinz (1997), which will be used to compare the degree of weathering of different samples at the same and different sites. Additionally, the proportion of *n*-alkanes and PAHs remaining through time will be compared among samples and sites. These analyses permit identification of the source of the oil.

LDPE data: The concentration and distribution of PAHs in these samples will be compared between oiled and non-oiled (control) armored beaches.

Boulder movement: We will use the same combination of survey methods employed in our earlier surveys. Measurement of boulder movement will be compared between years, by site. Various measures of movement, e.g., horizontal and vertical displacements, changes in angular orientation of the marker bolts, will be considered separately. Measurement error is determined through repeated measurements of selected marked bolts. The significance of displacements for the boulder armoring will be evaluated in relation to the size classes of the boulders on the beach. Variations between beaches will be contrasted, especially in relation to the extent of chemical weathering of oil samples.

D. Description of Study Area

As detailed above, we are proposing to continue monitoring of six sites located on the GOA coastline, in Katmai National Park & Preserve and Kenai Fjords National Park and Preserve (Irvine et al., 1999; 2006; 2007; Short et al., 2007). We have monitored oiling conditions and boulder movements at these sites since 1994. Maps with the location of the study sites and details of site morphology and sampling have been included in previous reports submitted to the EVOS Trustee Council.

E. Coordination and Collaboration with Other Efforts

NOAA is a cooperating agency, and Mark Carls, the head of the analytical lab at NOAA's Auke Bay Fisheries Laboratory, is a principal investigator on the project. We have been in communication with the NPS regarding this project, and most closely there with Bud Rice. We plan to have NPS staff with us in the field, and will be training staff in our sampling procedures. The NPS continues to be interested in and concerned with the persistence of oil on the Katmai and Kenai Fjords National Park coastlines.

III. SCHEDULE

A. Project Milestones

Objective 1. Determine status and extent of persistent oiling at the long-term GOA study sites.
To be met by March 2013

Objective 2. Determine if oil is leaking from GOA armored beaches.
To be met by March 2013

Objective 3. Determine the stability of the boulder armors.
To be met by February 2013

B. Measurable Project Tasks

FFY 11, 2nd quarter (January 1, 2011-March 31, 2011)
February: *Project funding approved by Trustee Council*

FFY 11, 3rd quarter (April 1, 2011-June 30, 2011)
Contracting, hiring, preparation for field work
Field work

FFY 11, 4th quarter (July 1, 2011-September 30, 2011)

Shipment of LDPE and mussel samples to Auke Bay Labs

FFY 12, 1st quarter (October 1, 2011-December 31, 2011)
December 15: *Begin hydrocarbon analyses of LDPEs and mussel samples*

FFY 12, 2nd quarter (January 1, 2012-March 31, 2012)

January 18: Annual Marine Science Symposium

March 1: Complete hydrocarbon analyses of LDPEs and mussel samples

Begin contracting for FY12 field work

FFY 12, 3rd quarter (April 1, 2012-June 30, 2012)

April 15: Submit annual report

Contracting, hiring, preparation for field work

FFY 12, 4th quarter (July 1, 2012-September 30, 2012)

Field work

Shipment of LDPE and hydrocarbon samples to Auke Bay Labs

Shipment of hydrocarbon samples to Woods Hole Oceanographic Institution

FFY 13, 1st quarter (October 1, 2012-December 31, 2012)

December 15: Begin data and hydrocarbon analyses

FFY 13, 2nd quarter (January 1, 2013-March 31, 2013)

January 18: Annual Marine Science Symposium

March 1: Complete hydrocarbon analyses

Write report/manuscript

FFY 13, 3rd quarter (April 1, 2013-June 30, 2013)

April 15: Submit final report to the Trustee Council Office. This will consist of a draft manuscript for publication.

FFY 13, 4th quarter (July 1, 2013-September 30, 2013)

Present findings at national conference (during FFY13 or FFY14)

C. Publications & Reports

The study results will be submitted to EVOS TC as a manuscript that will later be submitted for publication in a peer-reviewed journal. We are requesting funding for the writing of this manuscript and its publication in a peer-reviewed journal. The tentative title of one manuscript is: "Oil persistence 23-years after the *Exxon Valdez* spill on boulder-armored beaches distant from the spill origin." We plan to target the journal, *Marine Environmental Research*, with a submission date planned for Dec. 2013.

IV. LITERATURE CITED

Babcock, M.M., G.V. Irvine, P.M. Harris, J.A. Cusick, and S.D. Rice. 1996. Persistence of oiling in mussel beds three and four years after the *Exxon Valdez* oil spill. Pages 268-297 in S.D. Rice, R.B. Spies, D.A. Wolfe, and B.A. Wright, editors. Proceedings of the *Exxon Valdez* Oil Spill Symposium. American Fisheries Society Symposium 18.

Bishop, P. and Hughes, M. (1989). Imbricate and fitted fabrics in coastal boulder deposits on the Australian east coast. *Geology* 17, 544-547.

- Bodkin, J.L., B.E. Ballachey, T.A. Dean, A.K. Fukuyama, S.C. Jewett, L. McDonald, D.H. Monson, C.E. O'Clair, and G.R. VanBlaricom. 2002. Sea otter population status and the process of recovery from the 1989 '*Exxon Valdez*' oil spill. *Marine Ecology Progress Series* 241:237-253.
- Boufadel, M. C., Y. Sharifi, B. Van Aken, B. A. Wrenn, and K. Lee. 2010. Nutrient and oxygen concentrations within the sediments of an Alaskan beach polluted with the *Exxon Valdez* oil spill, *Environmental Science and Technology*, 44 (19), p 7418–7424.
- Carls, M.G., M.M. Babcock, P.M. Harris, G.V. Irvine, J.A. Cusick, and S.D. Rice. 2001. Persistence of oiling in mussel beds after the *Exxon Valdez* oil spill. *Marine Environmental Research* 51(2001): 167-190.
- Chapman, D. (2006). The virtual fish: SPMD basics. US Geological Survey Columbia River Research Center. wwwaux.cerc.cr.usgs.gov/SPMD/index.htm
- Dean, R.G. and Dalyrymple, R.A. (2004). *Coastal Processes with Engineering Applications*. Cambridge University Press, Cambridge.
- Esler, D. and S.A. Iverson. 2010. Female harlequin duck winter survival 11 to 14 years after the *Exxon Valdez* oil spill. *Journal of Wildlife Management* 74(3):471-478.
- Hayes, M.O. and Michel, J. (1999). Factors affecting the long-term persistence of *Exxon Valdez* oil in gravel beaches. *Marine Pollution Bulletin* 38, 92-101.
- Hayes, M.O., Michel, J., Betenbaugh, D. (2010). The intermittently exposed, coarse-grained gravel beaches of Prince William Sound, Alaska: comparison with open-ocean gravel beaches. *J. Coastal Research* 26(1): 4-30.
- Hills, E.S. (1970). Fitting, fretting and imprisoned boulders. *Nature* 226, 345-347.
- Irvine, G.V., Mann, D.H., Short, J.W. 1999. Multi-year persistence of oil mousse on high energy beaches distant from the *Exxon Valdez* spill origin. *Marine Pollution Bulletin* 38: 572-584.
- Irvine, G.V., Mann, D.H., Short, J.W. 2006. Persistence of ten-year old *Exxon Valdez* oil on Gulf of Alaska beaches: the importance of boulder armoring. *Marine Pollution Bulletin* 59(9): 1011-1022.
- Irvine, G.V., Mann, D.H., Short, J.W. 2007. Monitoring lingering oil from the *Exxon Valdez* spill on Gulf of Alaska armored beaches and mussel beds. Restoration Project 040708, Final Report, EVOS Trustee Council, Anchorage, AK.
- Li, H., M. C. Boufadel, Long-term persistence of oil from the *Exxon Valdez* spill in two-layer beaches, *Nature Geoscience*, 3, 96-99, 2010.

Michel, J. and Hayes, M.O. (1993a). Evaluation of the condition of Prince William Sound shorelines following the *Exxon Valdez* oil spill and subsequent shoreline treatment: Volume I: 1991 Geomorphological shoreline monitoring survey. National Oceanic and Atmospheric Administration Technical Memorandum NOS ORCA 67. 94 pp.

Michel, J. and Hayes, M.O. (1993b). Evaluation of the condition of Prince William Sound shorelines following the *Exxon Valdez* oil spill and subsequent shoreline treatment: Volume I: Summary of results - Geomorphological shoreline monitoring survey of the *Exxon Valdez* spill site, Prince William Sound, Alaska, Sept 1989 - Aug 1992. National Oceanic and Atmospheric Administration Technical Memorandum NOS ORCA 73.

Michel, J. and Hayes, M. O. 1999. Weathering patterns of oil residues eight years after the *Exxon Valdez* oil spill: Marine Pollution Bull., Vol. 38: 855-863.

Michel, J., Z. Nixon, M.O. Hayes, J. Short, G. Irvine, D. Betenbaugh, C. Boring, and D. Mann. 2010. Distribution of Subsurface Oil from the *Exxon Valdez* Oil Spill. Final Report, EVOS Restoration Project 070801.

Oak, H.L. (1984). The boulder beach: A fundamentally distinct sedimentary assemblage. Annals of the Association of American Geographers 74, 71-82.

Short, J.W., Lindeberg, M.R., Harris, P.M., Maselko, J.M., Pella, J.J. and Rice, S.D. 2004. Estimate of oil persisting on the beaches of Prince William Sound 12 years after the *Exxon Valdez* oil spill. Environmental Science and Technology. 38(1): 19-25.

Short, J.W., Irvine, G.V., Mann, D.H., Maselko, J.M., Pella, J.J., Lindeberg, M.R., Payne, J.R., Driskell, W.B., and Rice, S.D. 2007. Slightly weathered *Exxon Valdez* oil persists in Gulf of Alaska beach sediments after 16 years. Environmental Science and Technology 41:1245-1250.

Short, J.W., Maselko, J.M., Lindeberg, M.R., Harris, P.M., Rice, S.D. 2006. Vertical Distribution and Probability of Encountering Intertidal Exxon Valdez Oil on Shorelines of Three Embayments within Prince William Sound, Alaska. Environmental Science & Technology 40(12):3723-3729.

Vandermeulen, J.H. 1977. The Chedabucto Bay spill, *Arrow* 1970. Oceanus 20: 32-39.

Venosa, A.D., Campo, P., and Suidan, M.T. 2010. Biodegradability of lingering crude oil 19 years after the *Exxon Valdez* oil spill. Environmental Science and Technology 44(19):7613-7621.

Wentworth, C.K. (1922). A scale of grade and class terms for clastic sediments. Journal of Geology 30, 337-392.

Xia, Y, H. Li, M. C. Boufadel, and Y. Sharifi. 2010. Hydrodynamic factors affecting the persistence of the *Exxon Valdez* oil in a shallow bedrock beach, *Water Resources Research*, VOL. 46, W10528, 17 pp.

V. RESUMES

Gail V. Irvine

Research Ecologist
U.S. Geological Survey, Alaska Science Center
4210 University Drive
Anchorage, AK 99508

Phone: 907-786-7069
FAX: 907-786-7150
E-mail:girvine@usgs.gov

A. Education

Ph.D. 1983 University of California, Santa Barbara, Biological Sciences
Emphasis – Aquatic and Population Biology
M.S. 1973 University of Washington, Seattle; Zoology
B.A. 1969 University of California, Santa Barbara; Zoology (Honors)
Attended 1965-66 Alaska Methodist University

B. Professional Positions

Research Ecologist, U.S. Geological Survey, Alaska Science Center, 1995 – present
Coastal Resources Specialist, National Biological Survey/U.S. Geological Survey, 1993-1995
Coastal Resources Specialist, National Park Service, 1991-1993
Marine Biologist/Fisheries Scientist, Minerals Management Service, 1984-1991
Assistant Research Biologist, Marine Science Institute, University of California, Santa Barbara, 1983

C. Scientific Involvement

Editorial Board of Marine Systems Domain, online journal, TheScientificWorld, 2001-present
Member, Non-indigenous Species Working Group, PWS RCAC

D. Five Recent Publications Related to the Lingering Oil Project

- Irvine, G.V., Mann, D.H., Short, J.W. Sixteen-year persistence of slightly weathered *Exxon Valdez* oil under boulder armors on beaches distant from the 1989 spill. (*in revision*)
- Short, J.W., Irvine, G.V., Mann, D.H., Maselko, J.M., Pella, J.J., Lindeberg, M.R., Payne, J.R., Driskell, W.B., and Rice, S.D. 2007. Slightly weathered *Exxon Valdez* oil persists in Gulf of Alaska beach sediments after 16 years. *Environmental Science and Technology* 41:1245-1250.
- Irvine, G.V., D.H. Mann, and J.W. Short. 2006. Persistence of ten-year old *Exxon Valdez* oil on Gulf of Alaska beaches: The importance of boulder armoring. *Marine Pollution Bulletin* 59(9): 1011-1022.
- Irvine, G.V. 2000. Persistence of spilled oil on shores and its effects on biota. Chapter 126, pp. 267-281, *in* C.R.C. Sheppard, editor. *Seas at the Millennium: An Environmental Evaluation, Volume III, Global Issues and Processes*, Elsevier Science, Ltd., Oxford.

Irvine, G.V., D.H. Mann, J.W. Short. 1999. Multi-year persistence of oil mousse on high energy beaches distant from the *Exxon Valdez* spill origin. Marine Pollution Bulletin, 38(7): 572-584.

E. Other Publications

- Irvine, G. V. 2010. Development of monitoring protocols to detect change in rocky intertidal communities of Glacier Bay National Park and Preserve. USGS Open-file Report, 2010-1283.
- Pasch, A. D., N. R. Foster, and G. V. Irvine. 2010. Faunal analysis of late Pleistocene/early Holocene invertebrates provides evidence for paleoenvironments of a Gulf of Alaska shoreline inland of the present Bering Glacier margin, in Shuchman, R.A. and Josberger, E.G., eds., Bering Glacier: Interdisciplinary Studies of Earth's Largest Temperate Surging Glacier: Geological Society of America Special Paper 462, p. 251-274.
- Hallman, N. M. Burchell, B.R. Schoene, G.V. Irvine, and D. Maxwell. 2009. High-resolution sclerochronological analysis of the bivalve mollusk *Saxidomus gigantea* from Alaska and British Columbia: techniques for revealing environmental archives and archeological sensitivity. J. of Archeological Science 36: 2353-2364.
- Irvine, G.V. and E.N. Madison. 2008. Development of a monitoring protocol to detect ecological change in the intertidal zone of Sitka National Historical Park, Alaska. USGS Scientific Investigations Report 2008-5139.
- Carls, M.G., M.M. Babcock, P.M. Harris, G.V. Irvine, J.A. Cusick, and S.D. Rice. 2001. Persistence of oiling in mussel beds after the *Exxon Valdez* oil spill. Marine Environmental Research 51(2001): 167-190.

F. Persons with whom Dr. Irvine has collaborated on a project or publication with over the last four years

Scott Carpenter, Dept. of Geosciences, University of Iowa, Iowa City, Iowa

Sandra Lindstrom, Dept. of Botany, University of British Columbia, Vancouver, B.C., Canada

Dan Mann, Institute of Arctic Biology, University of Alaska, Fairbanks

Anne Pasch, Dept. Biology, University of Alaska, Anchorage

Jeanne Schaaf, Director, Lake Clark-Katmai Studies Center, National Park Service, Anchorage, AK

Nora Foster, NF Taxonomic Services, Fairbanks, Alaska

Jeff Short, Auke Bay Lab, NOAA, Juneau, Alaska

John Southon, Earth System Science Dept., University of California, Irvine, CA

Bernd Schoene, University of Mainz, Germany

Nadine Hallman, University of Mainz, Germany

Jacqui Michel, Research Planning, Inc., South Carolina

Zach Nixon, Research Planning, Inc., South Carolina

Miles Hayes, Research Planning, Inc., South Carolina

DANIEL H. MANN

CONTACT INFORMATION

email: dhmann@alaska.edu

mailing address: Scenarios Network for Alaska Planning, Geography Program, University of Alaska, Fairbanks, AK 99775

telephone: work (907) 474-7127

PROFESSIONAL CREDENTIALS

- 1983-1985 Post-Doctoral Fellow, Quaternary Research Center, University of Washington: Sea-level history and soil development in Svalbard.
- 1978-1983 PhD. Quaternary Science, College of Forest Resources, University of Washington: Quaternary History of the Lituya Glacial Refugium, Alaska.
- 1976-1978 MSc. Entomology, University of Washington: Ecology of Snowfield-Foraging Arthropods on Mount Rainier
- 1971-1975 B.A. Social Anthropology, University of Washington
-

EMPLOYMENT HISTORY

- 2008-present Associate Professor, Geography Program, School of Natural Resources University of Alaska, Fairbanks, Alaska
- 1994-2008 Research Associate, Institute of Arctic Biology, University of Alaska
- 1991-2006 Research Associate, Alaska Quaternary Center and Institute of Arctic Biology, University of Alaska
- 1988-1991 Research Associate, Quaternary Research Center, University of Washington
- 1985-1988 Director, School for Field Naturalists, University of Vermont
-

FIVE RELATED PUBLICATIONS

- Mann, D.H. and Streveler, G.P. (2008). Relative sea level history, isostasy, and glacial history in Icy Strait, Southeast Alaska. *Quaternary Research* 69, 201–216.
- Irvine, G.V., Mann, D.H., Short, J.W. Sixteen-year persistence of slightly weathered *Exxon Valdez* oil under boulder armors on beaches distant from the 1989 spill. (*in revision*)
- Short, J.W., Irvine, G.V., Mann, D.H., Maselko, J.M., Pella, J.J., Lindeberg, M.R., Payne, J.R., Driskell, W.B., and Rice, S.D. 2007. Slightly weathered *Exxon Valdez* oil persists in Gulf of Alaska beach sediments after 16 years. *Environmental Science and Technology* 41:1245-1250.
- Irvine, G.V., Mann, D.H., Short, J.W. 2006. Persistence of ten-year old *Exxon Valdez* oil on Gulf of Alaska beaches: the importance of boulder armoring. *Marine Pollution Bulletin* 59(9): 1011-1022.
- Irvine, G.V., D.H. Mann, J.W. Short. 1999. Multi-year persistence of oil mousse on high energy beaches distant from the *Exxon Valdez* spill origin. *Marine Pollution Bulletin*, 38(7): 572-584.

FIVE OTHER SIGNIFICANT PUBLICATIONS

- Mann, D.H., Groves, P., Reanier, R.E. and Kunz, M.L. (2010). Floodplains, cottonwood trees, hillslopes and peat: What happened the last time climate warmed suddenly in arctic Alaska? *Quaternary Science Reviews* 29, 3812-3830.
- Mann, D.H., Reanier, R.E., Beck, W., and Edwards, J. (2008). Drought, vegetation change, and human history on Rapa Nui (Isla de Pascua, Easter Island). *Quaternary Research* 69, 16-28
- Mann, D.H., Crowell, A.L., Hamilton, T.D., and Finney, B.P. (1999). Holocene Geologic and climatic history around the Gulf of Alaska. *Arctic Anthropology* 35, 112-131.
- Crowell, A.L. and Mann, D.H. (1996). Human populations, sea level change, and the archaeological record of the Northern Gulf of Alaska coastline. *Arctic Anthropology* 33, 16-37.
- Mann, D.H. and Hamilton, T.D. (1995). Late Pleistocene and Holocene Paleoenvironments of the North Pacific Coast. *Quaternary Science Reviews* 14, 449-471.

COLLABORATORS AND CO-AUTHORS (Last 4 years)

- Suzanne Anderson (Department of Geography, University of Colorado)
- James Edwards (Department of Surgery, Oregon Health Sciences University)
- John Edwards (Department of Zoology, University of Washington)
- Bruce Finney (Institute of Marine Science, University of Alaska)
- Pamela Groves (Institute of Arctic Biology, University of Alaska)
- Thomas Hamilton (US Geological Survey, Alaska Branch)
- David Meltzer (Department of Anthropology, Southern Methodist University)
- Scott Rupp (School of Natural Resources, University of Alaska)
- Ronald Sletten (Department of Geology, University of Washington)
- John Stone (Department of Geology, University of Washington)
- Gail Irvine (US Geological Survey, Alaska Science Center)
- Jeffrey Short (NOAA, Auke Bay Laboratory)
- Jacqui Michel (Research Planning, Inc., South Carolina)
- Zach Nixon (Research Planning, Inc., South Carolina)

Curriculum Vitae for Mark G. Carls

National Marine Fisheries Service, Auke Bay Laboratories
17109 Pt. Lena Loop Road
Juneau, AK 99801

email: mark.carls@noaa.gov
Phone: (907) 789-6019
FAX: (907) 789-6094

Education

M.Sc., 1978, biological oceanography, Dalhousie University, Halifax, Nova Scotia.
B.A., 1975, biology; Magna cum laude, Gustavus Adolphus College, St. Peter, Minnesota.
Additional coursework (30 semester hours), University of Alaska Southeast (statistics, genetics, fish, and misc)

Professional Experience

Fisheries Biologist, 1979-present, Auke Bay Laboratory.
Principal Investigator for *Exxon Valdez* Oil Spill Trustee Council

- Embryo toxicity: pink salmon, Pacific herring, zebrafish
- Pink salmon habitat
- Herring Synthesis
- Mussel and sediment contamination
- Hydrocarbon chemistry: sampling, interpretation, modeling

Biological Review Teams

- Pacific herring, Lynn Canal, Alaska (chairman)
- Status of Pacific herring in Puget Sound, Washington, Habitat and Ecological Processes Team

Recent related publications (lead author only)

Carls MG, Meador JP. 2010. A perspective on the toxicity of petrogenic PAHs to developing fish embryos related to environmental chemistry. *Human and Ecological Risk Assessment* 15:1084-1098.

Carls, M.G. 2006. Nonparametric identification of petrogenic and pyrogenic hydrocarbons in aquatic ecosystems. *Environ Sci Technol.* 40:4233-4239.

Carls, M.G., S.D. Rice, G.D. Marty, and D.K. Naydan. 2004. Pink salmon spawning habitat is recovering a decade after the *Exxon Valdez* oil spill. *Trans Am Fish Soc* 133:834-844.

Carls MG, LG Holland, JW Short, RA Heintz, SD Rice. 2004. Monitoring polynuclear aromatic hydrocarbons in aqueous environments with passive low-density polyethylene membrane devices. *Environ Toxicol Chem* 23:1416-1424.

Carls MG, Thomas RE, Rice SD. 2003. Mechanism for transport of oil-contaminated water into pink salmon redds. *Mar. Ecol. Prog. Ser.* 248:245-255.

Other significant publications

Carls MG, Thedinga J.F. 2010. Exposure of pink salmon embryos to dissolved polynuclear aromatic hydrocarbons delays development, prolonging vulnerability to mechanical damage. *Marine Ecology Research* 69:318-325

Carls MG, Holland L, Larsen M, Collier TK, Scholz NL, Incardona JP. 2008. Fish embryos are damaged by dissolved PAHs, not oil particles. *Aquatic Toxicology* 88:121-127.

Carls MG., PM Harris, SD Rice. 2004. Restoration of Oiled Mussel Beds in Prince William Sound, Alaska. *Mar. Environ. Res.* 57:359-376.

- Carls MG, GD Marty, JE Hose. 2002. Synthesis of the toxicological impacts of the *Exxon Valdez* oil spill on Pacific herring (*Clupea pallasi*) in Prince William Sound, Alaska, U.S.A. *Can. J. Fish. Aquat. Sci.* 59:1-20.
- Carls MG, MM Babcock, PM Harris, GV Irvine, JA Cusick, SD Rice. 2001. Mar Env Res 51:167-190.

Recent Collaborators (excluding ABL staff):

Dr. Mace Barron (P.E.A.K. Research)
Frederick C. Funk (consultant, Juneau, AK)
Dr. Jo Ellen Hose (Occidental University, CA)
Dr. John Incardona (NOAA, Northwest Fisheries Science Center)
Dr. John Kern (NOAA Damage Assessment and Restoration Center Northwest, Seattle, WA)
Dr. Gary Marty (Animal Health Centre, Abbotsford, BC)
Dr. Brenda Norcross (University of Alaska Fairbanks)
Dr. James Payne (Payne Environmental Consultants)
Dr. Terrance J. Quinn II (University of Alaska Fairbanks)
Dr. Robert Spies (Applied Marine Sciences, Livermore, CA)
Dr. Katherine Springman (University of California, Davis)
Dr. Bob Thomas (University of California, Chico).