

Project Title: Patterns and Processes of Population Change in Selected Nearshore Vertebrate Predators

Project Number: 02423
Restoration Category: Research and Monitoring
Proposers: Jim Bodkin, Dan Esler,
Brenda Ballachey
Lead Trustee Agency: DOI--USGS
Cooperating Agencies:
Alaska SeaLife Center: Yes
Project Duration: 4th year, 5-year project
Cost FY 02: \$329,700
Cost FY 03: \$250,000
Geographic Area: Prince William Sound
Injured Resource/Service: Sea Otter, Harlequin Duck

ABSTRACT

Sea otters and harlequin ducks have not fully recovered from the EVOS, based on population-level demographic differences between oiled and unoiled areas. Further, in oiled areas, both species show elevated cytochrome P4501A (CYP1A) through 1998, almost certainly reflecting continued exposure to oil. We propose to continue to explore links between oil exposure and the lack of population recovery, with the intent of understanding constraints to full recovery of these species and the nearshore environment generally. We also will monitor the progress of recovery of the species and the system. Proposed work consists of field components for both species, and a captive component for harlequin ducks. For sea otters, we will conduct aerial surveys of distribution and abundance and estimate age-specific survival rates. For harlequin ducks, field studies will examine the relationship between survival and CYP1A and, further, will serve to monitor these key parameters. Captive experiments on harlequin ducks will examine the relationships between oil exposure and CYP1A induction, and the metabolic and behavioral consequences of exposure to oil.

INTRODUCTION

The nearshore environment of Prince William Sound (PWS) received about 40% of the oil spilled after the *Exxon Valdez* ran aground (Galt et al. 1991). Concerns about nearshore recovery and restoration resulted in a suite of studies sponsored by the *Exxon Valdez* Oil Spill Trustee Council, including the Nearshore Vertebrate Predator project (NVP). Principal findings of NVP include an apparent lack of population recovery for sea otters (*Enhydra lutris*) and harlequin ducks (*Histrionicus histrionicus*), both invertebrate feeders in the nearshore ecosystem (Bodkin et al. 1999; Esler et al. 1999, Dean et al. 2000, Bodkin et al. in press). Over a three year period, harlequin ducks residing in oiled areas had poorer survival than those in unoiled areas (Esler et al. 2000a). Sea otters also experienced poor post-spill survival through 1998, based on modeling of ages-at-death (Monson et al. 2000). Further indication of increased mortality (or higher rates of emigration) of sea otters in oiled areas compared to their counterparts in unoiled areas is provided by inferences based on capture data (Bodkin et al. 1999, Bodkin et al. in press). Additionally, both species show evidence of continuing exposure to hydrocarbons, based on higher levels of the biomarker cytochrome P4501A (CYP1A), in oiled areas than unoiled (Ballachey et al. 1999). Elevations in CYP1A are not explained by background or natural hydrocarbon sources, as these were found to be negligible in intertidal areas of PWS (Short and Babcock 1996), nor by area differences in PCB contamination (Trust et al. 2000; USFWS unpub. data), leaving continued exposure to residual *Exxon Valdez* oil as the most plausible explanation. Residual oil is still stranded in intertidal areas of PWS (Babcock et al. 1996; Hayes and Michel 1999).

Conceptual links have been drawn describing mechanisms by which oil exposure could have population-level demographic impacts on sea otters and harlequin ducks. However, these links, and thus the processes that may limit full recovery, remain speculative. Therefore, we propose to build on the base of knowledge gained through previous research to (1) explore the relationships between oil exposure, individual health, and demographic attributes that could have population level effects, and (2) monitor the parameters identified in previous work that are effective and statistically powerful in describing population status and lend insight into the process of recovery of sea otters and harlequin ducks, and the nearshore environment generally.

Sea Otters

The NVP study provided several lines of evidence indicating that sea otters in the most heavily oiled portions of western Prince William Sound (WPWS), at northern Knight and Naked islands, have not recovered from oil-related injury (Bodkin et al. 1999, in press; Dean et al. 2000; Monson et al. 2000). The sea otter population at northern Knight has not increased between 1993-99 (the period for which we have aerial survey data), with numbers remaining at about half the estimated pre-spill abundance. Sea otters in oiled areas show reduced survival, relative to pre-spill rates (Bodkin et al. 1999; Monson et al. 2000). Levels of CYP1A were higher in sea otters from Knight Island than from unoiled reference areas, suggesting continued exposure to residual oil may be affecting recovery of the species. Additionally, increased proportions of larger-sized individuals of several sea otter prey species were identified at northern Knight, consistent with reduced predation and lack of recovery of the sea otter population in that area (Dean et al. 2000).

The sea otter component of this proposal builds on previous EVOS research (93045, 95025-99025) to develop a statistically sensitive and cost-effective program that will continue to track the WPWS sea otter population and nearshore ecosystem recovery, and investigate the effects of chronic oil exposure on sea otters. We will address the following questions: (1) are sea otters increasing in abundance in the most heavily oiled areas, and in western PWS overall ? and (2) has survival of sea otters returned to pre-spill rates?

Question 1 will be answered by continued aerial surveys of sea otter abundance at appropriate intervals to monitor the population and test predictions of a previously developed sea otter population model (Restoration study 99043; Udevitz et al. 1996). Surveys were done in 1999 and 2000, and will be conducted again in 2002. Sea otter aerial surveys were suspended in 2001, based on fiscal considerations and the need to re-sample sea otter survival and the bioindicator cytochrome P450 1A in that year. *This element is a continuation of work proposed and approved in Project 99423, and initiated in FY 1999.*

Question 2, regarding survival rates of sea otters, involves a modeling effort that utilizes ages-at-death of sea otters recovered as carcasses on beaches (Monson et al. 2000). This element was not initially included as part of Project 99423, but due to the compelling evidence of long-term injury provided by the modeling results in late 1999, the carcass surveys were added for FY2000-01 (supplementary funding provided in February 2000). We propose that carcass surveys be conducted again in 2002.

Harlequin Ducks

The most concerning result from NVP harlequin duck studies was the detection of significantly lower survival probabilities of adult females in oiled areas of PWS than in unoiled areas (Esler et al. 2000a). Analyses revealed that history of oil contamination was a more likely explanation for the survival difference than intrinsic differences between oiled and unoiled study areas. Further, projections of population trends using models incorporating these survival probabilities predicted declining populations on oiled areas and increasing populations on unoiled areas. This pattern was observed during Alaska Department of Fish and Game surveys (EVOSTC Project /427), suggesting that differences in survival were a likely mechanism for observed differences in population trends. Also, harlequin duck densities were lower on oiled Knight Island than on unoiled Montague Island, after accounting for intrinsic habitat differences; this is the pattern that would be predicted given high site fidelity and poorer survival on oiled areas. Finally, higher levels of CYP1A induction were detected on oiled areas.

Results from these recent studies lead to speculation that continued exposure to oil could result in poorer survival of harlequin ducks, which in turn would result in differences in population trends and densities. There are reasonable explanations for how oil may be related to survival (see Statement of Problem below). Unfortunately, however, these links are drawn from a wide array of sources, with limited inference to wild harlequin ducks in PWS. Thus, we propose studies that will explore the relationship between oil exposure and survival using both field and captive bird approaches. These will serve to examine mechanisms or processes that may continue to limit harlequin duck population recovery. These studies also will monitor the most critical elements revealed in previous studies to gauge the progress of recovery.

The specific questions that will be asked by the harlequin duck components of this study are: (1) what is the relationship between levels of oil exposure and CYP1A induction, and what levels of oil exposure result in CYP1A values similar to those measured in PWS? (2) are there metabolic or behavioral consequences of oil exposure that could be a mechanism by which harlequin duck survival is compromised? (3) is oil exposure (as indicated by CYP1A induction) related to survival of harlequin ducks in the wild? and (4) is contaminant exposure declining over time and, similarly, are survival rates on the oiled area improving through time? Questions 1 and 2 will be addressed using captive birds at the Alaska SeaLife Center during winters 2000-01 and 2001-02. Questions 3 and 4 will be addressed by biosampling and radio telemetry work during winters 2000-01, 2001-02, and 2002-03. *These studies are a continuation of work proposed and approved in Project 00423.* This work will examine both the process of recovery (through understanding of the mechanisms constraining population demography) and will monitor the progress of recovery by sampling survival and CYP1A induction of wild birds starting 3 years subsequent to the last work done as part of NVP (winter 1997-98).

NEED FOR THE PROJECT

A. Statement of Problem

Sea otters and harlequin ducks occupy an invertebrate-consuming trophic level in the nearshore and are conspicuous components of the nearshore ecosystem. In 1995, the NVP Project was initiated to examine the status of recovery of nearshore vertebrates (including sea otters, harlequin ducks, river otters and pigeon guillemots), and to evaluate possible causes for the apparent lack of recovery. Results of the NVP project clearly suggest that complete recovery has not occurred for sea otters and harlequin ducks, and the lack of recovery may be related to continued exposure to oil. This proposed work follows up on the critical elements revealed by the NVP studies.

Sea Otters

The sea otter population in WPWS was injured as a result of the spill. Estimates of sea otter mortality due to the spill range from 750 to 2,650 individuals (Garshelis 1997, Garrott et al. 1993). A population model (Udevitz et al. 1996) predicted recovery of the WPWS sea otter population in 10 to 23 years, projecting maximum annual growth rates from 0.10-0.14. Surveys to date (1993-1998) have shown a significant increasing trend in the WPWS sea otter population, averaging about 4% per year since 1993 (power > 0.80 to detect a 1% annual change in 5 annual WPWS surveys). In contrast to the western Sound overall, at northern Knight Island sea otter numbers remain below pre-spill estimates and do not show a significant increasing trend (Figure 1; Bodkin et al. 1999, Bodkin 2000; Dean et al. 2000).

Sea otter carcasses have been recovered from beaches in WPWS since 1976, thus providing one of the few long-term baseline data sets for evaluating post-spill injury. Carcass surveys initially were not proposed as part of Project 99423. However, in 1999 we applied recently developed modeling techniques (Doak and Morris 1999) to estimation of sea otter survival rates, utilizing

the distribution of otter ages-at-death as the basis for the model. The results provide compelling evidence of long-term injury from the EVOS (Monson et al. 2000). Briefly, the model involves a comparison of observed vs. predicted ages-at-death of sea otters prespill and postspill, using data from carcasses collected during 1976-98. Postspill survival of sea otters in the western Sound was poor relative to prespill rates, and by 1998, survival rates had not yet returned to prespill values. However, survival rates of younger age otters were increasing, suggesting that conditions were normalizing. These results are consistent with other observations of sea otters in western PWS, which suggest that the population in the most heavily oiled areas has not yet recovered (Figure 1). Carcass collections and modeling efforts based on age-at-death data may provide one of the most efficient tools for monitoring recovery of sea otters. Thus, we propose that carcass surveys (and subsequent modeling to estimate survival rates) be continued in 2002, as an additional tool for monitoring sea otter recovery in PWS.

The NVP study identified elevated expression of CYP1A in 6 species that inhabit the nearshore areas of WPWS, indicating continued exposure to residual EVOS oil (Ballachey et al. 1999). Sea otters were sampled in 1996-98, and in all years, animals from Knight and Naked islands (oiled area) had elevated CYP1A, compared to those from Montague Island (unoiled area; Figure 2). Further, levels at Montague were similar to those measured in otters from a relatively clean area in southeast Alaska with no known exposure to oil or other contaminants (USGS unpub. data). In 1998, the mean value of CYP1A in the oiled study area was lower than means for 1996 or 1997, suggesting exposure to residual oil is diminishing over time. We will resample the wild sea otter population for CYP1A in summer 2001 to determine if hydrocarbon exposure




Figure 1. Estimated sea otter abundance at northern Knight Island.

continues, and if so, if it has declined relative to levels measured in 1996-98. Sea otters in the most heavily oiled areas of WPWS will be targeted for sampling, with particular effort to capture those residing in the vicinities of known persistent oiled shoreline and bivalve populations (Hayes and Michel 1999, Fukuyama et al. in press) and oiled mussel beds (Harris et al. 2000), potentially enabling us to make a link between biomarker levels in sea otters and petroleum contaminants in mussels and sediments of their nearby habitat. Sea otters from Montague Island also will be captured to provide a non-exposed reference sample. We are not proposing further collection of sea otters until blood samples collected in 2001 are analyzed.




Figure 2. Measurement of cytochrome P4501A induction (RT-PCR technique) in sea otters in western Prince William Sound, 1996-98.

In summary, we propose continued monitoring of sea otter distribution, abundance, and survival rates in WPWS. These studies will be valuable in documenting actual recovery time for the nearshore system including sea otters, and providing long-term population trend data which may be used in assessing initial damage and subsequent recovery of sea otter populations in the event of future oil spills.

Harlequin Ducks

Harlequin ducks were, and remain, particularly vulnerable to deleterious effects of the oil spill. Much of the oil from the *Exxon Valdez* was deposited in the nearshore intertidal and shallow subtidal zones (Galt et al. 1991), the coastal habitats where harlequin ducks occur. Also, Goudie and Ankney (1986) suggested that harlequins were near the lower limit of body size for sea ducks occurring in environments similar to Prince William Sound in winter. Because harlequin ducks exist close to an energetic threshold, any perturbation (e.g., an oil spill) that either affects health or condition directly (via toxic effects or increased metabolic costs) or indirectly (via food abundance) could have significant consequences for the population.

Also, among ducks, sea duck life histories are particularly K-selected (Eadie et al. 1988). Harlequin ducks typically defer reproduction for 3 years, have relatively low annual investment in reproduction, and are long-lived (Goudie et al. 1994). Species with these characteristics have relatively low potential rates of population change and, thus, following a perturbation such as an oil spill, require many years in the absence of continued adverse effects to recover to previous population levels. Further, population dynamics of animals with this life history strategy are particularly sensitive to variation in adult survival (Goudie et al. 1994, Schmutz et al. 1997).

Sea ducks have a general pattern of high philopatry throughout their annual cycle (e.g., Limpert 1980, Savard and Eadie 1989) and harlequin ducks follow this pattern, having high fidelity to

molting and wintering sites (Robertson 1997; Esler, unpubl. data). High site fidelity could result in vulnerability to population effects because: (1) if residual oil spill damages exist, birds from oiled areas are vulnerable to spill effects as they return to those areas annually (i.e., these birds are affected disproportionately and are subject to cumulative effects), and (2) if dispersal and movements among areas are limited, recovery of groups of birds in oiled areas can occur only through demographic processes specific to that group (i.e., numbers are not enhanced through immigration from other areas). High site fidelity is an adaptive behavioral strategy in natural situations and predictable environments (Robertson 1997), but does not accommodate movement to undisturbed sites in the face of human-caused perturbations.

Evidence from recent studies (NVP and /427) suggests that, as might be predicted from their vulnerability, harlequin duck populations have not fully recovered and, in fact, continue to suffer deleterious effects from the oil spill. Over the course of 3 winters, survival probabilities differed between oiled and unoiled areas (Figure 3). Survival probabilities were high, and similar between areas, in fall. However, survival diverged between areas during mid-winter, presumably the period during which conditions are most difficult for harlequin ducks. Also, differences in CYP1A induction were detected between populations from oiled and unoiled areas (Figure 4; Trust et al. 2000), although this was measured on different birds than those for which survival data were collected. Further, body mass during winter showed a slight, negative relationship with CYP1A level.

One can speculate on mechanisms by which continued exposure to oil could be related to differences in survival probabilities. Most lab studies have shown that mallards are tolerant of internal ingestion of oil, with toxic effects not evident until very high doses. These studies have been used to suggest that harlequin ducks should, similarly, be unaffected by residual Exxon Valdez oil (Stubblefield et al. 1995, Boehm et al. 1996). However, other studies have found that, with addition of other stressors




Figure 3. Survival probabilities of harlequin ducks.




Figure 4. Comparison of CYP1A induction (hepatic EROD activity) in harlequin ducks from Prince William Sound.

such as cold temperatures, oiled ducks in the lab suffered considerably higher mortality than unoiled (Holmes et al. 1978, 1979). This seems to be a much more appropriate analog for wild harlequin ducks. Particularly given their vulnerability to spill effects and hypothesized existence near an energetic threshold, harlequin ducks may not be able to handle additive effects of the oil spill, even if relatively small.

To fully understand the process of harlequin duck population recovery from the oil spill, it is important to address these speculated links between oil exposure and survival probabilities, and subsequently population trends. The research proposed here is designed to explore these potential mechanisms constraining population recovery through field studies of winter survival and CYP1A induction and captive studies of metabolic, behavioral and CYP1A responses to controlled oil exposure. Further, because of their susceptibility to spill effects and high site fidelity, harlequin ducks are an ideal species for monitoring recovery of the nearshore environment.

B. Rationale/Link to Restoration

Sea otter and harlequin duck restoration requires assessments of population recovery status and definition of impediments to recovery. For harlequins and sea otters, the proposed work incorporates monitoring activities which, given the “baseline” data collected in NVP and other post-spill studies, will allow us to gauge recovery status. Additionally, the research components proposed herein represent a comprehensive approach to understanding the factors that affect population dynamics and definition of critical bottlenecks to recovery. Without an understanding of the underlying processes that dictate population change, we can not prescribe specific activities to enhance recovery. The project directly addresses the restoration objectives both by examining the processes affecting recovery and by monitoring the progress of recovery, including survival rates and contaminant exposure.

Sea Otters

Recovery of sea otters will be complete when population size returns to estimated pre-spill abundance, and there is no further evidence of continuing exposure to residual oil. Sea otter restoration requires an understanding of population status and the processes affecting changes in population status. Continued monitoring of sea otter distribution, abundance, survival rates and prey populations in WPWS will provide insight into recovery and improve future recovery models, and potentially allow us to document the actual recovery time for the nearshore system, including sea otters. A further benefit of these project components is provision of long-term population trend data and monitoring tools which may be used in assessing initial damage and subsequent recovery of sea otter populations in the event of future oil spills.

Harlequin Ducks

Harlequin duck restoration will be complete when densities have recovered to prespill levels and birds no longer show evidence of oil contamination. Poor survival in oiled areas is the most plausible cause for lack of recovery to prespill densities; restoration requires an understanding of the factors that affect survival rates, in particular the effects of oil exposure. The restoration

objectives for harlequin ducks are addressed both by examining the processes affecting recovery and by monitoring the progress of recovery, in particular contaminant exposure.

C. Location

Studies will be conducted in PWS. Specific study sites for the sea otter components will be northern Knight Island and Port Chalmers/Stockdale at Montague Island, as used in the NVP project. Harlequin duck study sites also will be those used in previous NVP work: unoiled Montague Island and oiled Green Island, Crafton Island, Main Bay and Foul Bay. Captive studies will be done at the Alaska SeaLife Center in Seward. Communities affected by the project include Chenega, Whittier, Cordova and Seward.

COMMUNITY INVOLVEMENT AND TRADITIONAL ECOLOGICAL KNOWLEDGE

The project will continue to inform and coordinate our community involvement activities, including the collection of indigenous knowledge with Dr. Henry Huntington, TEK specialist Chugach Regional Resources Commission and Hugh Short, Community Coordinator, EVOS Restoration Office. We will continue to solicit advice from the above parties and gather information on TEK through local community facilitators and residents. Efforts have and will continue to be made throughout the restoration process to participate in and provide public involvement in the design and implementation of this project. Information gathered from this project will be shared with local communities. Project staff has and will continue to present information to local communities or prepare articles or photographs for Trustee Council publications. Boat and air charter contracts, and other services will be contracted from local sources when possible.

PROJECT DESIGN

A. Objectives

Sea Otters

Field Studies

1. Estimate of sea otter abundance and population trends over time in WPWS overall, and in oiled and unoiled study areas within WPWS.
2. Monitor progress of sea otter population recovery via tracking of survival rates in oiled areas.

Harlequin Ducks

Field Studies

1. Estimate winter survival rates of harlequin ducks in relation to area (history of oil contamination) and indices of oil exposure (CYP1A induction).
2. Monitor progress of harlequin duck population recovery via tracking of survival rates and CYP1A induction in oiled and unoiled areas.

Captive Studies

1. Measure the CYP1A response in oil-exposed, captive harlequin ducks.
2. Quantify the metabolic and behavioral consequences of oil exposure.

B. Methods

The proposed research employs field studies on sea otters, and both field studies and experimental work with harlequin ducks. This combination of approaches addresses the need for controlled work to look explicitly at the effects of oil exposure on hypothesized mechanisms of mortality and field work to document the relevance of those mechanisms under wild conditions. With captive studies on harlequin ducks, we propose to quantify metabolic and behavioral responses to known regimes of oil exposure as well as indicate the level of oil exposure that corresponds to CYP1A induction detected in the field. For both species, field studies are necessary to understand the relevance of these relationships to animals in the wild, and to monitor population and system recovery.

Sea Otters

Field Studies

The proposed sea otter work employs aerial surveys to track population abundance and growth. This approach will provide information on recovery status of the population, assessed through trends in population size. Additional components proposed for 2002 are collection of carcasses for determination of ages at death, to be used in estimation of age-specific survival rates.

Sea otter population monitoring--We will continue to use previously developed aerial survey techniques which employ counts along systematic transects, and intensive search units (ISU's) to estimate a correction factor for each survey (Bodkin and Udevitz, 1999). We will conduct a single survey of the entire WPWS in 2002, and in alternate years, conduct a survey of the entire PWS. From the combination, we will obtain an estimated population size for WPWS annually (except in 2001). We will continue annual replicate surveys (5 or more replications per survey) of the smaller NVP study sites, initiated in 1993 (except in 2001).

Carcass surveys--Age specific survival estimates will be generated based on age distributions of the dying portion of the population, evaluated through recovery of beach-cast sea otter carcasses in western PWS. Beaches will be surveyed once during late April or early May after snow melt

but prior to summer revegetation, which may hide carcasses washed high on the beach by winter storms. Data recorded for each carcass include: (1) relative location of carcass on the beach, (2) relative condition and completeness of carcass, (3) position of remains relative to previous year's vegetation, (4) relative age (adult, subadult, pup), (5) sex, and (6) specimens collected (e.g., entire carcass, skull, baculum, none). Skulls (when present) will be taken from all carcasses and a tooth extracted for aging (Bodkin et al. 1997). Any fresh carcasses collected will be necropsied as soon as possible and tissue samples collected for potential toxicology and histopathology studies.

Harlequin Ducks

Field Studies

The key data for field studies are paired CYP1A and survival data, which will allow for explicit tests of the hypothesis that mortality and oil exposure are related in wild harlequin ducks. We intend to collect survival and exposure data from 50 birds in each of 3 years by capturing them during early winter, conducting surgeries to both implant transmitters and biopsy livers, and monitoring subsequent winter survival. These types of data have been successfully collected during NVP studies.

This research requires capture of flighted harlequin ducks during early winter, after they have been on wintering sites long enough to be potentially exposed to residual oil, yet before the mid-winter period when survival probabilities diverged during NVP studies (Figure 3). The mid-winter period is presumably the time of greatest stress and thus the period when oil spill effects would be most likely to be expressed as differences in survival probabilities. The interval between capture and the critical mid-winter period must allow for at least a 2-week censor period to ensure that survival data are not biased by effects of capture, handling, or surgery (Esler et al. 2000b; Mulcahy and Esler 1999). Thus, we propose capturing birds during a 3-week period in November to generate both survival data and exposure data from the same individuals.

We will use floating mist nets (Kaiser et al. 1995) to catch flying birds in oiled (Knight Island, Green Island, Crafton Island, Main Bay, Foul Bay) and unoiled (Montague Island) study areas. Use of the same study areas as the NVP project allows for direct comparisons of results. The floating mist net capture technique was used successfully during NVP studies. However, this technique does not allow handling of as many birds as molt drives, so age cohorts used in survival estimation will not be as restricted as in NVP studies, which included only after-third-year females. We will radio females of all age classes; age parameters will be included in all analyses to account for any survival differences due to these effects. Captured birds will be banded with uniquely coded USFWS bands, aged by bursal probing (Mather and Esler 1999), and sexed by plumage characteristics.

To estimate survival probabilities of harlequin ducks, we will use implantable radio transmitters with external antennas (Korschgen et al. 1996). Implanted transmitters have been successfully used in waterfowl studies (e.g., Olsen et al. 1992, Haramis et al. 1993), and an increasing body of literature suggests that radio transmitters implanted into wild waterfowl are less disruptive than external methods of attachment, based on differences in survival or return rates (Ward and

Flint 1995, Dzus and Clark 1996), behavior (Pietz et al. 1993), and reproductive rates (Pietz et al. 1993, Rotella et al. 1993, Ward and Flint 1995, Paquette et al. 1997), especially for diving ducks (Korschgen et al. 1984). NVP studies (Esler et al. 2000b) demonstrated that recapture probabilities of radio-marked harlequin ducks were not lower than unradioed individuals. Surgeries will be conducted by certified veterinarians experienced in avian implant surgeries, following procedures outlined in Alaska Biological Science Center, USGS Biological Resources Division standard protocol. Transmitters will weigh approximately 18g, which is < 3% of the body mass of the smallest wintering female harlequin ducks captured during NVP studies. Transmitters will be equipped with mortality sensors; the pulse rate will change from 45 to 90 beats per minute when a mortality is indicated. Mortality status will be confirmed by either carcass recovery or detection of signals from upland habitats, which are not used by harlequin ducks during nonbreeding periods.

We will conduct radio telemetry flights at approximately weekly intervals from the capture and marking period through the end of March. Survival data entry and general description will follow procedures outlined in Pollock et al. (1989a, 1989b), as modified by Bunck et al. (1995). We will examine effects of area, season, and CYP 1A on survival by comparing AIC_c values (Burnham and Anderson 1998) among models with different combinations of these effects. The AIC_c indicates the most parsimonious model by balancing the goodness-of-fit of each model (from the maximum likelihood) with the number of parameters to be estimated. Under this approach, the model with the lowest AIC_c indicates the combination of parameters that are best supported by the data, which we will interpret as the factors related to variation in survival. Survival estimates and variances will be calculated by iterative solution of the likelihood using program MARK (White and Burnham 1999).

CYP1A induction will be measured by EROD activity. Small liver biopsies (approximately 0.1 g) will be surgically removed and immediately frozen in a liquid nitrogen shipper. EROD activity analyses will be conducted in a contracted lab following standard procedures (Trust et al. 2000). Plumage swabs (Duffy et al. 1999) and plucked feathers will be used to assess presence of external oil.

Captive Studies

Captive bird studies will examine metabolic, behavioral, and biomarker responses to known oil-exposure regimes. This work is designed to experimentally test effects of oil exposure on parameters that are hypothesized to influence dynamics of wild harlequin duck populations; these effects are impossible to assess under field conditions.

Harlequin ducks to be used in captive studies will be captured during wing molt from unoiled parts of PWS. During molt, harlequin ducks congregate and are susceptible to capture by herding flocks of flightless birds into pens (Clarkson and Goudie 1994). Birds will be banded with USFWS bands and with individually coded plastic tarsus bands. Tarsus bands will be oriented to be read from bottom to top as the bird is standing. Sex will be identified based on plumage characteristics and age class determined by bursal probing (Mather and Esler 1999). Body mass of all birds at capture will be measured.

Following capture, 25 females older than second-year will be flown to the Alaska SeaLife Center in Seward. We will use 21 birds for the winter 2001-02 experiments; we will release any birds that are not adapting well to captivity or are not needed for experiments back to capture sites. Captured individuals will undergo quarantine and adjustment periods prior to any experimental manipulation or oil exposure. Captive birds will be housed in outdoor pens to expose them to natural climatic and photoperiod conditions. Oil exposure will be designed to simulate long-term, intermittent exposure, which is likely similar to exposure experienced by wild birds. The experimental design for winter 2001-02 experiments will be determined following analyses of data collected during the first year of the experiment, which recently ended (March 2001). Oil exposure will continue through the critical mid-winter period and behavioral and metabolic measures will be taken throughout the winter. Because CYP1A sampling requires a liver biopsy, we will get only 1 measure of induction, taken in late winter. Following a 2-week post-surgery recovery period (without any exposure), captive birds will be released in the area of their original capture.

Behavior of captive birds will be quantified using time-activity observations throughout winter for all exposure levels. Behavioral categories will follow those used in studies of wild harlequin ducks (Goudie and Ankney 1986, Fischer 1998), e.g., feeding, resting, swimming, courtship, etc. Time-activity budgets will be contrasted among treatments.

Metabolic consequences of oil exposure will be quantified using two approaches: doubly-labeled water to estimate daily energy expenditure (DEE) and oxygen consumption to estimate basal metabolic rate (BMR). This approach will allow different views into the metabolic effects of exposure. DEE is a measure of existence costs over longer (1-3 day) time periods. DEE incorporates all of the metabolic costs during this time; elevated DEE in exposed birds would be consistent with a hypothesis of oil exposure increasing existence costs with potential survival implications. Similar DEE among treatments but different activity levels (see above) also would have implications for survival under natural conditions. BMR estimates metabolism without costs of thermoregulation, digestion, and activity; these data will assess whether background metabolic costs are higher in exposed than unexposed birds. Body mass of all individuals also will be measured at all handling events; these data will be interpreted in light of metabolic and behavioral measurements.

DEE estimation using doubly-labeled water requires injection of water with both the oxygen and water isotopically-labeled. As the hydrogen is lost only through water and oxygen through both water loss and carbon dioxide production, the difference in turnover rates between marked hydrogen and oxygen can be used to estimate metabolism. BMR will be measured using a flow-through respirometer to measure oxygen consumption. An oxygen analyzer is on site at the Alaska SeaLife Center and was used in winter 2000-01 experiments. BMR of all birds will be measured throughout the winter, including prior to any exposure to establish background rates.

CYP1A induction of all captive birds will be measured at the end of the experiment by EROD activity, described above. EROD activity will be compared among all treatments.

C. Cooperating Agencies, Contracts, and Other Agency Assistance

USGS-BRD personnel will be responsible for directing and conducting sea otter and harlequin duck studies.

Contract with Dr. Dan Esler for the harlequin duck components.

SCHEDULE

A. Measurable Project Tasks for FY02

Sea Otters

December-March: Coordinate and plan aerial surveys, carcass collections, sea otter capture, community involvement, prepare equipment.
Obtain/update marine mammal permits.

April-May: Collection of beach-cast sea otter carcasses for survival estimates.

July: Aerial surveys of sea otters in PWS

August - April Data analysis and report prep

Harlequin Ducks

Oct-March: Conduct studies of captive flock at the Alaska SeaLife Center, with birds captured during late FY01.

November: Capture harlequin ducks for field studies of survival and CYP1A induction.

Nov-March: Monitor radioed birds for survival study.

March: Surgically biopsy livers of captive birds for EROD activity; after a recovery period, birds will be released at the original capture site.

April - August: Prepare for field studies (e.g., order radios, contact boat charter operators, maintain winter trap, contact biosample contractors, etc.).

B. Project Milestones and Endpoints

This is a projected five-year research and monitoring program (initiated FY99, with completion of all objectives by FY03; see below) designed to assess the recovery of two injured species. Project objectives will be assessed annually. At the end of each year results will be compared with the restoration goals to assess whether recovery has occurred. The reporting schedule is described below, and is consistent with EVOS Trustee Council guidelines.

Sea Otters

- FY01-02: Field studies (aerial and carcass surveys) are scheduled to occur from April through July, 2002.
- FY03: Complete analysis, prepare final report and manuscript preparation

Harlequin Ducks

- FY01-03: Captive bird experimental work is scheduled for winters 2000-01 and 2001-02. Field studies are scheduled to occur from November through March, winters 2000-01, and 2001-02; occurrence of field work in winter 2002-03 will depend on review of harlequin duck population recovery status in 2002.

C. Completion Date

All project objectives will be met by FY03.

PUBLICATIONS AND REPORTS

Annual reports will be presented to the Chief Scientist by April 15. An annual report of FY02 activities will be submitted to the Restoration Office on or before 15 April 2003. A final report will be prepared at the end of the proposed work unless continued monitoring is warranted or when recovery objectives are met. Special reports (publications) will be prepared during the course of the study if warranted. Publications will be prepared for peer-review journals when sufficient data have been collected.

PROFESSIONAL CONFERENCES

D. Esler attendance at 2002 American Ornithologists Union meeting, date and location to be determined. J. Bodkin attendance at biennial Conference on Biology of Marine Mammals, November 2001, Vancouver, BC.

NORMAL AGENCY MANAGEMENT

The work proposed here is not part of normal agency management and is related specifically to research addressing oil spill restoration concerns. No similar work has been conducted, is currently being conducted, or is planned using agency funds.

COORDINATION AND INTEGRATION OF RESTORATION EFFORT

As described in the Introduction, this research relies on incorporation of data from other Trustee sponsored research, including projects /025 and /427. Equipment and commodities purchased under /025 will be used to conduct the proposed research and data collection and analysis will follow previously established protocols and standards.

EXPLANATION OF CHANGES IN CONTINUING PROJECTS

In 1998, the EVOS Trustee Council first approved funding for Restoration Project 99423, “Patterns and Processes of Population Change in Sea Otters”, an extension of the NVP project. The objectives of the project included sea otter aerial surveys of PWS, replicate surveys of sea otters at Knight and Montague Islands and sampling of sea urchin populations. In 1999, the Trustee Council approved the addition of harlequin duck studies to 00423 with the revised project title “Patterns and Processes of Change in Selected Nearshore Vertebrates”. Those studies included relating harlequin survival to oil exposure and captive studies to assess responses to controlled oil exposure. In February 2000, the Trustee Council approved an amendment to 00423, to fund carcass recovery surveys in WPWS, to collect data on sea otter ages at death for estimation of survival rates.

In July 2000, the project 01423 budget and DPD were revised to reflect suspension of the aerial surveys for sea otters in July 2001. Because salary costs were included in aerial surveys and also supported urchin work that is discontinued in 2001, salary costs of 28.8 K were redirected from aerial surveys to sea otter biomarker and survival sampling.

PROPOSED PRINCIPAL INVESTIGATORS

James Bodkin
Alaska Biological Science Center
USGS-Biological Resources Division
1011 E. Tudor Rd., Anchorage, Alaska 99503
PHONE: (907) 786-3550
FAX: (907) 786-3636
james_bodkin@usgs.gov

Dan Esler
Centre for Wildlife Ecology
Simon Fraser University
5421 Robertson Road, RR1
Delta, BC V4K 3N2 Canada
PHONE: (604) 940-4652
FAX: (604) 946-7022
desler@sfu.ca

Brenda Ballachey
Alaska Biological Science Center
USGS-Biological Resources Division
1011 E. Tudor Rd., Anchorage, Alaska 99503
PHONE: (907) 786-3512
FAX: (907) 786-3636
brenda_ballachey@usgs.gov

PRINCIPAL INVESTIGATOR QUALIFICATIONS

Jim Bodkin, Research Wildlife Biologist, and team leader for coastal ecosystem in Alaska for the Alaska Biological Science Center of USGS, Biological Resources Division. He has over 40 peer-reviewed scientific publications and directs an active coastal marine research program. He has studied and published on sea otter foraging ecology and community structuring since 1988 and has been principal investigator for sea otter survey methods development. He earned a M.S. from California State Polytechnic University in 1986.

Dan Esler is a University Research Associate with Simon Fraser University in British Columbia. He has conducted waterfowl research in arctic and subarctic regions of Alaska and Russia for the past 12 years. Since 1995 he has served as project leader for harlequin duck studies as part of the EVOSTC-sponsored Nearshore Vertebrate Predator project. He earned a M.S. from Texas A&M University in 1988 and a Ph.D. from Oregon State University in 2000. He has authored over 20 peer-reviewed journal publications and numerous reports and presentations addressing research and issues in waterbird conservation.

Brenda Ballachey is a Research Physiologist at the Alaska Biological Science Center of USGS, Biological Resources Division. She was Project Leader for sea otter NRDA studies from 1990 through 1996, and has been involved in all aspects of post-spill research on sea otters, including the Nearshore Vertebrate Predator (NVP) project, with primary responsibilities for examining effects of residual oil on biomarkers and health of sea otters and other NVP study species. She received her M.S. in 1980 at Colorado State University, and Ph.D. in 1985 Oregon State University. She has authored or coauthored over 25 peer-reviewed publications.

KEY COOPERATORS

Sea Otters

Daniel Monson
Alaska Biological Science Center
USGS-Biological Resources Division
1011 E. Tudor Rd.
Anchorage, AK 99503
PHONE: (907) 786-3449
FAX: (907) 786-3636
daniel_monson@usgs.gov

Paul W. Snyder, DVM, PhD.
Department of Veterinary Pathobiology
Purdue University
West Lafayette, IN 470907-1243
PHONE: (765) 494-9676
FAX: (765) 494-9830
pws@vet.purdue.edu

Harlequin Ducks

Kim Trust
U. S. Fish and Wildlife Service
Ecological Services
605 W. 4th Avenue, Room G62
Anchorage, AK 99501
(907) 271-2783
kim_trust@mail.fws.gov

Daniel Mulcahy, DVM, PhD.
Alaska Biological Science Center
USGS-Biological Resources Division
1011 E. Tudor Rd.
Anchorage, AK 99503
(907) 786-3451
daniel_mulcahy@usgs.gov

Dan Rizzolo
Dept. of Fisheries and Wildlife
Oregon State University
Corvallis, OR 97331
(541) 737-1969
rizzolo@mailbox.orst.edu

LITERATURE CITED

- Babcock, M.M., G.V. Irvine, P.M. Harris, J.A. Cusick, and S.D. Rice. 1996. Persistence of oiling in mussel beds three and four years after the *Exxon Valdez* oil spill. *Am. Fish. Soc. Symp.* 18:286-297.
- Ballachey, B.E., J.L. Bodkin, S. Howlin, K.A. Kloecker, D.H. Monson, A.H. Rebar and P.W. Snyder. 1999a. Appendix BIO-01 in NVP Draft Final Report (Project 950 Hematology and serum chemistry of sea otters in oiled and unoled areas of Prince William Sound, Alaska, from 1996-98. 25-99025).
- Ballachey, B.E., J.J. Stegeman, P.W. Snyder, G.M. Blundell, J.L. Bodkin, T.A. Dean, L. Duffy, D. Esler, G. Golet, S. Jewett, L. Holland-Bartels, A.H. Rebar, P.A. Seiser, and K.A. Trust. 1999b. Oil exposure and health of nearshore vertebrate predators in Prince William Sound following the *Exxon Valdez* oil spill. Chapter 2 in NVP Draft Final Report (Project 95025-99025).
- Bodkin, J.L. and M.S. Udevitz. 1999. An aerial survey method to estimate sea otter abundance. In: Garner GW, Amstrup SC, Laake JL, Manly BJF, McDonald LL, Robertson DG (eds) *Marine mammal survey and assessment methods*. AA Balkema, Rotterdam, p 13-26.
- Bodkin, J.L., J.A. Ames, R.J. Jameson, A.M. Johnson, and G.E. Matson. 1997. Estimating age of sea otters with cementum layers in the first premolar. *J. Wild. Manage.* 61(3):967-973.

- Bodkin, J.L., B.E. Ballachey, T.A. Dean, S. Jewett, L. McDonald, D. Monson, C. O'Clair, and G. VanBlaricom. 1999. Recovery of sea otters in Prince William Sound following the *Exxon Valdez* oil spill. Chapter 3A in NVP Draft Final Report (Project 95025-99025).
- Bodkin, J.L., B.E. Ballachey, T.A. Dean, A. K. Fukuyama, S. Jewett, L. McDonald, D. Monson, C. O'Clair, and G. VanBlaricom. In press. Sea otter population status and the process of recovery from the 1989 Exxon Valdez oil spill. Marine Ecology Progress Series.
- Boehm, P. D., P. J. Mankiewicz, R. Hartung, J. M. Neff, D. S. Page, E. S. Gilfillan, J. E. O'Reilly, and K. R. Parker. 1996. Characterization of mussel beds with residual oil and the risk to foraging wildlife 4 years after the *Exxon Valdez* oil spill. *Env. Toxicol. and Chem.* 15:1289-1303.
- Bunck, C. M., C-L. Chen, and K. H. Pollock. 1995. Robustness of survival estimates from radio-telemetry studies with uncertain relocation of individuals. *J. Wildl. Manage.* 59:790-794.
- Burnham, K. P., and D. R. Anderson. 1998. Model selection and inference: a practical information theoretic approach. Springer-Verlag, New York, New York, USA.
- Clarkson, P., and R.I. Goudie. 1994. Capture techniques and 1993 banding results for moulting harlequin ducks in the Strait of Georgia, B.C. Pages 11-14 in Proc. 2nd Harlequin Duck Symp., Hornby Island, B.C.
- Dean, T.A., J.L. Bodkin, A.K. Fukuyama, S.C. Jewett, D.H. Monson, C.E. O'Clair, and G.R. VanBlaricom. 2001. Food limitation and the recovery of sea otters in Prince William Sound. Marine Ecology Progress Series. In press.
- Dean, T.A., J.L. Bodkin, S.C. Jewett, D.H. Monson and D. Jung. 2001. Changes in sea urchins and kelp following a reduction in sea otter density as a result of the *Exxon Valdez* oil spill. Marine Ecology Progress Series. In press.
- Doak, D.F. and W.F. Morris. 1999. Detecting population-level consequences of ongoing environmental change without long-term monitoring. *Ecology* 80:1537-1551.
- Duffy, L. K., M. K. Hecker, G. M. Blundell, and R. T. Bowyer. 1999. An analysis of the fur of river otters in Prince William Sound, Alaska: oil related hydrocarbons 8 years after the *Exxon Valdez* oil spill. *Polar Biol.* 21:56-58.
- Duggins, D.O. 1980. Kelp beds and sea otters: an experimental approach. *Ecology* 61:447-453.
- Dzus, E. H., and R. G. Clark. 1996. Effects of harness-style and abdominally implanted transmitters on survival and return rates of mallards. *J. Field Ornith.* 67:549-557.
- Eadie, J. M., F. P. Kehoe, and T. D. Nudds. 1988. Pre-hatch and post-hatch brood amalgamation in north American Anatidae: a review of hypotheses. *Can. J. Zool.* 66:1709-1721.
- Esler, D., T.D. Bowman, K.A. Trust, B.E. Ballachey, T.A. Dean, S. Jewett, and C. O'Clair. 1999. Harlequin duck perspective: Mechanisms of impact and potential recovery of nearshore vertebrate predators. Chapter 4 in NVP Draft Final Report (Project 95025-99025).
- Esler, D., J. A. Schmutz, R. L. Jarvis, and D. M. Mulcahy. 2000a. Winter survival of adult female harlequin ducks in relation to history of contamination by the Exxon Valdez oil spill. *Journal of Wildlife Management* 64: in press.
- Esler, D., D. M. Mulcahy, and R. L. Jarvis. 2000b. Testing assumptions for unbiased estimation of survival of radio-marked harlequin ducks. *Journal of Wildlife Management* 64:591-598.

- Estes, J.A. and J. F. Palmisano. 1974. Sea otters: their role in structuring nearshore communities. *Science* 185:1058-1060.
- Estes, J.A., N.S. Smith and J.F. Palmisano. 1978. Sea Otter Predation and Community Organization in the Western Aleutian Islands, Alaska. *Ecology* 59(4):822-833.
- Estes, J. A. and D. O. Duggins. 1995. Sea otters and kelp forests in Alaska: generality and variation in a community ecological paradigm. *Ecological Monographs* 65(1):75-100.
- Exxon Valdez* Oil Spill Trustee Council. 1996. *Exxon Valdez* Oil Spill Restoration Plan. Draft Update on Injured Resources & Services. Anchorage.
- Fischer, J. B. 1998. Feeding behavior, body condition, and oil contamination of wintering harlequin ducks at Shemya Island, Alaska. M.S. Thesis, Univ. of Mass., Amherst.
- Fukuyama, A.K., G. Shigenaka and R.Z. Hoff. In press. Effects of residual *Exxon Valdez* oil on intertidal *Prototheca staminea*: mortality, growth and bioaccumulation of hydrocarbons in transplanted clams. *Mar. Poll. Bull.*
- Galt, J. A., W. J. Lehr, and D. L. Payton. 1991. Fate and transport of the *Exxon Valdez* oil spill. *Environ. Sci. Technol.* 25:202-209.
- Garrott, R.A., L.L. Eberhardt and D.M. Burns. 1993. Mortality of sea otters in Prince William Sound following the *Exxon Valdez* oil spill. *Mar. Mam. Sci.* 9:343-359.
- Garshelis, D. L. 1997. Sea otter mortality estimated from carcasses collected after the Exxon Valdez oil spill. *Conservation Biology.* 11(4):905-916.
- Goudie, R. I., and C. D. Ankney. 1986. Body size, activity budgets, and diets of sea ducks wintering in Newfoundland. *Ecology* 67:1475-1482.
- Goudie, R. I., S. Brault, B. Conant, A. V. Kondratyev, M. R. Petersen, and K. Vermeer. 1994. The status of sea ducks in the North Pacific rim: toward their conservation and management. *Proc. North Am. Wildl. and Nat. Res. Conf.* 59:27-49.
- Haramis, G. M., D. G. Jorde, and C. M. Bunck. 1993. Survival of hatching-year female canvasbacks wintering in Chesapeake Bay. *J. Wildl. Manage.* 57:763-771.
- Harris, P., M. Carls, and C. Brodersen. 2000. Monitoring of oiled mussel beds in Prince William Sound (abstract). 2000 Restoration Workshop, January 18-19, 2000. EVOS Trustee Council, Anchorage.
- Hayes, M.O. and J. Michel. 1999. Factors determining the long-term persistence of Exxon Valdez oil in gravel beaches. *Marine Pollution Bulletin* 38(2):92-101.
- Holland-Bartels, L. et al. 1997. Mechanisms of impact and potential recovery of nearshore vertebrate predators. Exxon Valdez Oil spill restoration project annual report 96025. April, 1997.
- Holland-Bartels, L. et al. 1998. Mechanisms of impact and potential recovery of nearshore vertebrate predators. Exxon Valdez Oil spill restoration project annual report 97025. April, 1998.
- Holmes, W. N., J. Cronshaw, and J. Gorsline. 1978. Some effects of ingested petroleum on seawater-adapted ducks (*Anas platyrhynchos*). *Env. Res.* 17:177-190.
- Holmes, W. N., J. Gorsline, and J. Cronshaw. 1979. Effects of mild cold stress on the survival of seawater-adapted mallard ducks (*Anas platyrhynchos*) maintained on food contaminated with petroleum. *Env. Res.* 20:425-444.
- Johnson, A. M. 1987. Sea otters of Prince William Sound, Alaska. U.S. Fish and Wildlife Service, Alaska Fish and Wildlife Research Center, Unpublished Report. 86pp.
- Kaiser, G. W., A. E. Derocher, S. Crawford, M. J. Gill, and I. A. Manley. 1995. A capture technique for marbled murrelets in coastal inlets. *J. Field Ornithol.* 66:321-333.

- Korschgen, C. E., S. J. Maxson, and V. B. Kuechle. 1984. Evaluation of implanted radio transmitters in ducks. *J. Wildl. Manage.* 48:982-987.
- Korschgen, C. E., K. P. Kenow, A. Gendron-Fitzpatrick, W. L. Green, and F. J. Dein. 1996. Implanting intra-abdominal radio transmitters with external whip antennas in ducks. *J. Wildl. Manage.* 60:132-137.
- Lebreton, J. D., and J. Clobert. 1991. Bird population dynamics, management, and conservation: the role of mathematical modeling. Pages 105-125 in Perrins, C.M., J. D. Lebreton, and G. J. M. Hirons (eds.). *Bird population studies: relevance to conservation and management.* Oxford Univ. Press.
- Limpert, R. J. 1980. Homing success of adult buffleheads to a Maryland wintering site. *J. Wildl. Manage.* 44:905-908.
- Mather, D. D., and D. Esler. 1999. Evaluation of bursal depth as an indicator of age class of harlequin ducks. *Journal of Field Ornithology* 70:200-205.
- Monson, D.H., D.F. Doak, B.E. Ballachey, A. Johnson, and J.L. Bodkin. 2000. Long-term impacts of the *Exxon Valdez* oil spill on sea otters, assessed through age-dependent mortality patterns. *Proc. Nat'l. Acad. Sciences, USA* 97(12):6562-6567.
- Mulcahy, D. M., and D. Esler. 1999. Surgical and immediate postrelease mortality of harlequin ducks implanted with abdominal radio transmitters with percutaneous antennae. *Journal of Zoo and Wildlife Medicine* 30:397-401.
- Mulcahy, D. M., D. Esler, and M. K. Stoskopf. 1999. Loss from harlequin ducks of abdominally implanted radio transmitters equipped with percutaneous antennas. *Journal of Field Ornithology* 70:244-250.
- Olsen, G. H., F. J. Dein, G. M. Haramis, and D. G. Jorde. 1992. Implanting radio transmitters in wintering canvasbacks. *J. Wildl. Manage.* 56:325-328.
- Paquette, G. A., J. H. Devries, R. B. Emery, D. W. Howerter, B. L. Joynt, and T. P. Sankowski. 1997. Effects of transmitters on reproduction and survival of wild mallards. *J. Wildl. Manage.* 61:953-961.
- Pietz, P. J., G. L. Krapu, R. J. Greenwood, and J. T. Lokemoen. 1993. Effects of harness transmitters on behavior and reproduction of wild mallards. *J. Wildl. Manage.* 57:696-703.
- Pollock, K. H., S. R. Winterstein, C. M. Bunck, and P. D. Curtis. 1989a. Survival analysis in telemetry studies: the staggered entry design. *J. Wildl. Manage.* 53:7-15.
- Pollock, K. H., S. R. Winterstein, and M. J. Conroy. 1989b. Estimation and analysis of survival distributions for radio-tagged animals. *Biometrics* 45:99-109.
- Reidman, M.L. and J.A. Estes. 1990. The sea otter (*Enhydra lutris*): Behavior, ecology and natural history. Biological Report 90(14). U.S. Fish and Wildlife Service, Washington, D.C. 126 pp.
- Robertson, G.J. 1997. Pair formation, mating system, and winter philopatry in harlequin ducks. PhD dissertation, Simon Fraser University, Vancouver, B.C.
- Rotella, J. J., D. W. Howerter, T. P. Sankowski, and J. H. Devries. 1993. Nesting effort by wild mallards with 3 types of radio transmitters. *J. Wildl. Manage.* 57:690-695.
- Savard, J-P. L., and J. McA. Eadie. 1989. Survival and breeding philopatry in Barrow's and common goldeneyes. *Condor* 91:198-203.
- Short, J. W., and M. M. Babcock. 1996. Prespill and postspill concentrations of hydrocarbons in mussels and sediments in Prince William Sound. *Am. Fish. Soc. Symp.* 18:149-166.

- Schmutz, J. A., R. F. Rockwell, and M. R. Petersen. 1997. Relative effects of survival and reproduction on population dynamics of emperor geese. *J. Wildl. Manage.* 61:191-201.
- Snyder, P.W., T. Kondratyuk, B.E. Ballachey and J. Vanden Heuvel. 1999. CYP1A gene expression in sea otters (*Enhydra lutris*): a quantitative reverse transcriptase-polymerase chain reaction to measure CYP1A mRNA in peripheral blood mononuclear cells. Appendix BIO-02 *in* NVP Draft Final Report (Project 95025-99025).
- Stubblefield, W. A., G. A. Hancock, W. H. Ford, and R. K. Ringer. 1995. Acute and subchronic toxicity of naturally weathered *Exxon Valdez* crude oil in mallards and ferrets. *Env. Toxicol. and Chem.* 14:1941-1950.
- Trust, K. A., D. Esler, B. R. Woodin, and J. J. Stegeman. 2000. Cytochrome P450 1A induction in sea ducks inhabiting nearshore areas of Prince William Sound, Alaska. *Marine Pollution Bulletin* 40:397-403.
- Udevitz, M.S., B.E. Ballachey and D.L. Bruden. 1996. A population model for sea otters in western Prince William Sound. Exxon Valdez oil spill restoration project final report (restoration project 93043-3), National Biological Service, Anchorage, AK. 34pp.
- Vanden Heuvel, J.P., G.C. Clark, C.L. Thompson, Z. McCoy, C.R. Miller, G.W. Lucier, and D.A. Bell. 1993. CYP1A mRNA levels as a human exposure biomarker: use of quantitative polymerase chain reaction to measure CYP1A expression in human peripheral blood lymphocytes. *Carcinogenesis* 14(10):2003-2006.
- Vanden Heuvel, J.P., G.C. Clark, M.C. Kohn, A.M. Tritscher, W.F. Greenlee, G.W. Lucier, and D.A. Bell. 1994. Dioxin-responsive genes: Examination of dose-response relationships using quantitative reverse transcriptase-polymerase chain reaction. *Cancer Research* 54:62-68.
- Ward, D. H., and P. L. Flint. 1995. Effects of harness-attached transmitters on premigration and reproduction of brant. *J. Wildl. Manage.* 59:39-46.
- White, G. C., and K. P. Burnham. 1999. Program MARK -- survival estimation from populations of marked animals. *Bird Study* 46 Supplement:120-138.

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

Budget Category:	Authorized FY 2001	Proposed FY 2002	PROPOSED FY 2002 TRUSTEE AGENCIES TOTALS					
			ADEC	ADF&G	ADNR	USFS	DOI	NOAA
							\$329.7	
Personnel	\$169.0	\$66.2						
Travel	\$13.7	\$3.9						
Contractual	\$145.3	\$223.5						
Commodities	\$29.2	\$10.5						
Equipment	\$1.0	\$0.0	LONG RANGE FUNDING REQUIREMENTS					
Subtotal	\$358.2	\$304.1				Estimated FY 2003		
General Administration	\$35.5	\$25.6						
Project Total	\$393.7	\$329.7				\$250.0		
Full-time Equivalents (FTE)	0.0	1.0						
	Dollar amounts are shown in thousands of dollars.							
Other Resources	\$0.0	\$0.0			\$0.0	\$0.0		
Comments:								

FY02

Prepared: 7/5/01

Project Number: 02423
 Project Title: Pattern and Process of Population Change in
 Selected Nearshore Vertebrates
 Lead Agency: DOI--USGS

FORM 2A
 MULTI-
 TRUSTEE

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

Budget Category:	Authorized FY 2001	Proposed FY 2002						
Personnel	\$169.0	\$55.7						
Travel	\$13.7	\$3.9						
Contractual	\$145.3	\$223.5						
Commodities	\$29.2	\$10.5						
Equipment	\$1.0	\$0.0	LONG RANGE FUNDING REQUIREMENTS					
Subtotal	\$358.2	\$293.6			Estimated			
General Administration	\$35.5	\$24.0			FY 2003			
Project Total	\$393.7	\$317.6			\$250.0			
Full-time Equivalents (FTE)		0.9						
Dollar amounts are shown in thousands of dollars.								
Other Resources								
Comments:								

FY02

Prepared: 7/5/01

Project Number: 02423
 Project Title: Pattern and Process of Population Change in
 Selected Nearshore Vertebrates
 Agency: DOI--USGS

FORM 3A
 TRUSTEE
 AGENCY

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

Contractual Costs:		Proposed
Description		FY 2002
Charter vessel, sea otter beach walks - 10 days @ \$1200 (so)		12.0
OAS aerial survey costs 80 hr @ 220/hr (so)		17.6
Matson's Laboratory - tooth ages, 75 @ \$5 (so)		0.4
4A Linkage #2 Simon Fraser University (hd)		161.5
Oregon State Univerisity - graduate student support (hd)		32.0
Contractual Total		\$223.5
Commodities Costs:		Proposed
Description		FY 2002
Misc field/office supplies (so)		3.0
Equipment maintenance and repair (so)		1.0
Fuel (so)		2.0
Vet supplies (hd)		4.5
Commodities Total		\$10.5

FY02

Project Number: 02423
 Project Title: Pattern and Process of Population Change in
 Selected Nearshore Vertebrates
 Agency: DOI--USGS

FORM 3B
 Contractual &
 Commodities

Prepared: 7/5/01

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

Budget Category:	Authorized FY 2001	Proposed FY 2002						
Personnel	\$0.0	\$10.5						
Travel	\$0.0	\$0.0						
Contractual	\$0.0	\$0.0						
Commodities	\$0.0	\$0.0						
Equipment	\$0.0	\$0.0	LONG RANGE FUNDING REQUIREMENTS					
Subtotal	\$0.0	\$10.5			Estimated FY 2003			
General Administration	\$0.0	\$1.6						
Project Total	\$0.0	\$12.1			\$0.0			
Full-time Equivalents (FTE)		0.1						
Dollar amounts are shown in thousands of dollars.								
Other Resources								
Comments:								

FY02

Prepared: 7/5/01

Project Number: 02423
 Project Title: Pattern and Process of Population Change in
 Selected Nearshore Vertebrates
 Agency: DOI--FWS

FORM 3A
 TRUSTEE
 AGENCY

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

Contractual Costs:		Proposed
Description		FY 2002
		0.0
		0.0
		0.0
		0.0
		0.0
Contractual Total		\$0.0
Commodities Costs:		Proposed
Description		FY 2002
		0.0
		0.0
		0.0
		0.0
		0.0
Commodities Total		\$0.0

FY02

Prepared: 7/5/01

Project Number: 02423
 Project Title: Pattern and Process of Population Change in Selected Nearshore Vertebrates
 Agency: DOI--FWS

FORM 3B
 Contractual &
 Commodities

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

Budget Category:	Authorized FY 2001	Proposed FY 2002						
Personnel		\$69.0						
Travel		\$6.4						
Contractual		\$68.5						
Commodities		\$17.6						
Equipment		\$0.0						
Subtotal	\$0.0	\$161.5	LONG RANGE FUNDING REQUIREMENTS					
Indirect		\$0.0			Estimated FY 2003			
Project Total	\$0.0	\$161.5						
Full-time Equivalents (FTE)		1.0						
Dollar amounts are shown in thousands of dollars.								
Other Resources								
<p>Comments: SIMON FRASER UNIVERSITY</p> <p>No overhead or fees are charged by the university on this contract.</p>								

FY02

Prepared: 7/5/01

Project Number: 02423
 Project Title: Pattern and Process of Population Change in Selected Nearshore Vertebrates
 Agency: DOI-USGS --Simon Fraser University Contract

FORM 4A
 Non-Trustee
 SUMMARY

2002 EXXON VALDEZ TRUSTEE COUNCIL PROJECT BUDGET

October 1, 2001 - September 30, 2002

New Equipment Purchases:		Number of Units	Unit Price	Proposed FY 2002
Description				
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
				0.0
Those purchases associated with replacement equipment should be indicated by placement of an			New Equipment Total	\$0.0
Existing Equipment Usage:		Number of Units		
Description				

FY02

Prepared: 7/5/01

Project Number: 02423
 Project Title: Pattern and Process of Population Change in Selected Nearshore Vertebrates
 Agency: USGS--Simon Fraser University Contract

FORM 4B
 Equipment
 DETAIL