

Exxon Valdez Oil Spill
Restoration Project Final Report

Prince William Sound Food Webs: Structure and Change

Restoration Project 01393
Final Report

Thomas C. Kline, Jr., Ph. D. (Principal Investigator)

Signe Baumann (Laboratory Technician)

Prince William Sound Science Center
P. O. Box 705
Cordova, Alaska 99574

April 2002

Prince William Sound Food Webs: Structure and Change

Restoration Project 01393

Final Report

Study History: Project 393 commenced as project 99393 in April 2000 and is ending as project 01393 in April 2002. Years 1 and 2 focused on supplementing the isotopic database to resolve Ecopath model validation data gaps. The results of this project goal were presented in the 2001 Annual Report. A portion of these results were published as two Wakefield symposium volumes. The second goal was a retrospective isotopic analysis based on the periostracum of *Mytilus californianus*. Preliminary results reported in the 2001 annual report suggested modifications to sampling methods. These changes were carried out during the final project year, which are reported herein.

Abstract: Project 393 was funded to address gaps in our knowledge of Prince William Sound ecological processes based on recent stable isotope. In particular, stable isotope research has shown that the advective regime connecting the northern Gulf of Alaska (GOA) with Prince William Sound (PWS) fluctuates in its effect upon nutritional processes in Fishes on an annual basis. This project: (1) conducted a retrospective analysis carbon-13 since EVOS, and (2) addressed Ecopath model validation data gaps. These analyses may enable a better understanding of the ecological role of ‘regime shift’ processes conjectured to be impeding the natural restoration of populations in PWS affected by the EVOS.

Key Words: Ecosystem, Food Webs, Prince William Sound, Stable Isotopes

Project Data: *Description of data* - Data consist of natural carbon and nitrogen stable isotope abundance measurements expressed in intentionally recognized delta units. *Format* - Data are being published in the scientific manuscript in the form of tables and figures. *Custodian* - Contact Dr. Thomas C. Kline, Jr., Prince William Sound Science Center, P. O. Box 705, Cordova, AK 99754. e-mail: tkline@pwssc.gen.ak.us. *Availability* – in the open scientific literature or contact Dr. Kline for specific data, which can be e-mailed.

Citation: Kline, T. C., Jr. and S. Baumann 2002. Prince William Sound Food Webs: structure and change, *Exxon Valdez* Oil Spill Restoration Project Final Report (Restoration Project 01393), Prince William Sound Science Center, Cordova, Alaska.

Table of Contents

Introduction	4
Figure 1	6
Figure 2	7
Methods	8
Results	10
Table 1.	10
Figure 3	11
Figure 4	15
Literature Cited	18

List of Tables

Table 1. Summary comparing whole soft tissues with periostracum mean stable isotope values.

List of Figures

Figure 1. Inter-decadal shift in $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ content of bowhead whale baleen; from Schell (2001).

Figure 2. Seasonal shifts in the in $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ content of bulk net zooplankton, from Kline (1999).

Figure 3. $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of *Mytilus californianus* periostracum samples.

Figure 4. Fractional growth-year isotopic patterns.

Published Results

Kline, T.C., Jr. 2001. The trophic position of Pacific herring in Prince William Sound Alaska based on their stable isotope abundance. *In*: F. Funk, J. Blackburn, D. Hay, A.J. Paul, R. Stephenson, R. Toresen, and D. Witherell (eds.), Herring: Expectations for a New Millennium. University of Alaska Sea Grant, AK-SG-01-04, Fairbanks. p. 69-80.

Kline, T. C. Jr. 2002. Relative Trophic Position of *Cancer magister* megalopae. *In*: (eds.) Crabs in Cold Water Regions: Biology, Management, and Economics. University of Alaska Sea Grant, AK-SG-02-01, Fairbanks. p.645-649.

Executive Summary

Recent research has shown that the oceanographic conditions connecting the northern Gulf of Alaska with Prince William Sound may affect nutritional processes in fishes. Accordingly, food webs are subject to changes in carbon flow occurring between Gulf of Alaska and Prince William Sound. This project sought to (a) conduct retrospective analysis of Gulf of Alaska production shifts since the oil spill, and (b) address Ecopath model validation data gaps. Under direction of Chief Scientist objective (b) was deleted during the second project year. Nevertheless, two manuscripts derived from that aspect

of this study were published during 2001, re-prints of which are included as appendices. The report narrative, however, addresses objective (a).

Introduction

Shifts in carbon flow, which were driven by variations in the physical environment, represent fundamental changes in the way the PWS ecosystem supports commercially important species and is a probable mechanism explaining regime shifts. PWS fish populations may thus depend upon physical processes that regulate the availability of their macro-zooplankton forage. Stable isotope analysis (SIA) suggested that a portion of variability of macro-zooplankton may be ascribed to a varying subsidy of oceanic sources that are transported by currents into PWS from the adjacent Gulf of Alaska (Kline 1999).

The SIA approach, which uses stable isotope ratios of carbon and nitrogen, is unique in its ability to integrate time and spatial scales at mesoscale levels. The tracer aspect of the SIA is analogous to artificial tracer experiments without the burden of needing to generate signals or experimental artifacts. Because SIA provided the evidence suggesting that the Gulf carbon subsidy of PWS pelagic production appears to vary among years (Kline 2001), it is a useful GEM parameter for resolving how oceanographic processes affect fisheries recruitment over longer time scales. For example, SIA suggested long-term shifts from retrospective analysis of bowhead whale baleen (Figure 1; Schell 2001). Long-term systematic shifts among a collection of baleen from several whales provide evidence for change although the nature of that change is not yet understood (Schell, 2001).

Furthermore, the shifts observed in baleen exceed those predicted by the Seuss effect (which was expected to be notable, given the multi-decade nature of the data series in the baleen study) and thus reflect local environmental changes (Schell 2001). Investigation into the nature of isotopic change was therefore needed. There is also substantial seasonal isotopic variability (Figure 2; Kline 1999). Seasonal affects accounted for up to 40% of the variability in SIA of net plankton from one growth period. It is not know whether this pattern exists every year. The large intra-annual (seasonal) and inter-annual (e.g., Kline 2001, Schell 2001) may also provide clues for the observed long-term shifts. However, at present there is no long-term context for the Gulf of Alaska and PWS as provided by whale baleen for the Bering Sea.

The presently available limited temporal extent of the SIA database for PWS suggested a need for expansion through retrospective analysis. Shelled mollusks secrete a protein outer covering while they grow. In the case of bivalves the oldest material is located on the umbone, typically the pointed end of the shell, while the youngest is at the lip, where the mantle edge is in contact with the shell. The mantle secretes the protein layer, periostracum, along with CaCO₃-conatining parts of the shell. Winter growth checks enable aging the shell and the layer of periostracum above it. Mytilid bivalves have a particularly thick periostracum layer that spontaneously separates from the shell as it dries. Periostracum samples can thus be aged from from animals collected live. The objective of this project was thus to perform SIA on the periostracum of *Mytilus*

californianus and assess its potential paleoindicator for the Northern Gulf of Alaska. For the purposes of this study it was assumed that once laid down, periostracum is metabolically inert, thus preserving changes that took place at the time it was laid down. Evidence that this assumption comes from the fact that once damaged, periostracum is not replaced (in mussels), effecting 'bald spots.'




Figure 1. Inter-decadal shift in $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ content (vertical axis) of bowhead whale baleen ascribed to feeding in the Bering Sea; from Schell (2001).




Figure 2. Seasonal shifts in the in $^{15}\text{N}/^{14}\text{N}$ (A) and $^{13}\text{C}/^{12}\text{C}$ (B (raw data) & C (lipid-normalized)) content (vertical axis) of bulk net zooplankton collected during March to June 1995 (Julian Day = Day of the Year) in the northern Gulf of Alaska and Prince William Sound, from Kline (1999). Panels B and C reflect increase in correlation resulting from correction for lipid content.

Methods

Sea-mussels, *Mytilus californianus*, were collected at Middleton Island in September 1997 with help from Cordova fishermen. They provided the Principal Investigator with the knowledge and opportunity to acquire the *Mytilus californianus* samples that are providing an approach to retrospective analysis in the present study. Middleton Island's location in the Alaska Current provides an "upstream perspective" on the EVOS area since samples from there reflect changes in plankton upstream before interaction with PWS-origin carbon is possible. *Mytilus californianus* were collected from a rocky reef exposed at low tide (~ 0.5m MLLW) from the southeast shore of Middleton Island on 16 September 1997. From a collection measuring approximately 50 L in volume (three large buckets), 50 *Mytilus californianus* with mostly intact periostracum (showing little wear) and showing little to no signs of bio-fouling were "high-graded" and saved frozen (-20 °C). Stable isotopic analysis (SIA) was made on the outer protein layer (periostracum) on the shells and body tissues of *Mytilus californianus* of varying ages from this sub-collection.

Periostracum was initially sampled at the annulus level for isotopic analyses (Kline 2001). Growth checks were scored using a scalpel. Mussels were placed in a freeze-drier. Freeze-drying caused the periostracum to separate from the calcareous portion of the mussel shell. Flakes of periostracum that were peeling off the shell were forcep'd from each annulus and placed in a vial. Each periostracum sample (~ 2 mg) was sub-sampled according to the need of the then newly installed Finnegan Delta Plus stable isotope ratio mass spectrometer at the UAF Stable Isotope Facility. The new machine facilitated making several SIA from each annulus. However, the SD of the data was large (Kline 2001), suggesting the need for a more refined periostracum sampling technique. Developing and implementing micro periostracum sampling thus became the goal of project 393 during 2001-2. The problem consisted of collecting, sequentially, a series of small pieces of periostracum from along the long-axis of the shell. The difficulty lay in controlling the spontaneous peeling or flaking of periostracum material. Several methods were attempted – the rubber cement band technique, however, was successful and is described below.

Whole (soft tissues intact) mussel specimens were stored frozen (-20 °C) until preparation commenced. A scalpel was used to make two parallel incisions, approximately 10 mm apart, along the major axis of a shell valve while still frozen. Incisions ran from the narrow or beak end to the edge of the valve most distal to the beak. The incisions were re-traced several times to ensure that the resulting grooves were as deep as possible. A few additional narrow incisions were made within the two main tracks to facilitate sample collection. The mussels were placed in a freeze dryer (Labconco Shelf Drier) at -30 °C, which was operated for ~ 1-2 hours following evacuation. Freeze-drying was observed during this period to mitigate unwanted spontaneous separation of the periostracum from the valve. Mussels were removed from the freeze-dryer at the first

signs of separation or peeling. Rubber cement has the property of self-adhesion in preference to the object it is attached to, minimizing contamination of the object. Approximately 10 to 20% of the material in a sample would have to be contaminant to significantly affect isotopic determination, with less effect occurring with smaller differences in the isotopic composition between the contaminant and sample. A layer of rubber cement (Stanford) was then applied between the two incisions and allowed to air dry, which took ~ 30-40 minutes. Mussels were then placed back into freeze dryer set at - 30 °C and re-evacuated. The temperature setting of the freeze drier was gradually increased to + 10 °C over a three day period. During this time, rubber cement was applied twice daily, re-evacuating each time, until temperature of freeze dryer reached about +10 degrees. The freeze-drying process was completed with the freeze dryer set at 20 °C for 24 hours. Mussels were then removed from the freeze-drier. The rubber cement was gently lifted from the mussel while a scalpel was run underneath to ensure a clean break of the attached and intact periostracum strip. The strip, rubber cement side down, was tacked onto a styrofoam block. The block facilitated holding the periostracum while ~1-3 mm long sections were removed while being examined under a stereo microscope (Wild) set at low power. A scalpel and fine-tip forceps were used to handle the small samples, which ranged in length from 1.3mm-2.8mm, average 2.25mm. The samples, which were ~ 0.3 mg dry mass each, were collected from the valve's edge or lip towards the beak and placed into separate vials. The beginning and ending position of each sampled relative to the valve lip was measured with a dial micrometer. The soft tissues of each mussel were saved in a vial after being removed from the shell and ground to a fine powder.

Annuli positions were assessed on mussel valves after removal of the periostracum and dried soft tissues. A growth check was ascertained to be an annulus and not a 'false check' if it formed a continuous perimeter line. Annuli generally co-occurred with a change in slope of the valve surface that caused a shadow when examined under direct oblique sunlight (< 30 ° elevation) using a low power stereo microscope (Wild). The distance of each annulus was measured using a dial micrometer to the nearest 0.1 mm.

Samples were sent to the UAF Stable Isotope Facility for analysis using a Finnegan Delta Plus stable isotope ratio mass spectrometers equipped with Conflo II and III front-ends. Stable nitrogen and carbon isotope ratios, $^{15}\text{N}/^{14}\text{N}$ and $^{13}\text{C}/^{12}\text{C}$ are reported using conventional delta notation reflecting the natural abundance deviation from international standards, air N_2 and VPDB.

Isotope data analysis was facilitated on a Macintosh G4 micro-computer and the Statview (SAS, Inc.) analysis software. This software was used to plot the data according to mean sample position for each mussel. The mean integer value for each isotope among all samples was used as a y-axis reference value for plotting annuli positions with isotope data, which were placed on the x axis. Seasonal isotopic analysis was facilitated by converting sample positions (spatial) to fractional growth-year (implied temporal) such that the starting annulus position had a value of 0 while the ending position was 1. Data

plotted according to fractional growth year were limited to those years for which a series of data (generally ≥ 3 observations) were available. Cubic spline fits to the data were used. The resulting plots were assessed qualitatively

Effects of calcium carbonate contamination on $^{13}\text{C}/^{12}\text{C}$ values were estimated to increase values systematically by 1.3 based on the difference of acid and non-acid cleaned periostracum from a pool of five mussels. Samples were not acid cleaned to avoid $^{15}\text{N}/^{14}\text{N}$ artifacts.



Results

The three mussels prepared for SIA using the rubber cement band technique are summarized in Table 1. The Stable isotope data versus valve position and annuli position for mussels A, B and C are shown in Figure 3. These figures include the integer values nearest the mean $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ were +8 and -18, respectively, in conjunction with the indicated annuli positions.



Table 1. Summary comparing whole soft tissues with periostracum mean stable isotope values.

Mussel ID	Length of Shell (mm)	Date processed	Total length of periostracum sample	Sample Increment (mm)	Number samples taken	Number of data points	Soft tissues N-15	Soft tissues C-13 (raw data)	Periostracum N-15	Periostracum N-15 <i>SD</i>	Periostracum C-13	Periostracum C-13 <i>SD</i>
A	115	Oct-01	66	2	34	31	8.8	-20.2	7.8	0.4	-18.2	1.0
B	104	Oct-01	72	3	25	21	9.1	-18.7	8.1	0.5	-17.9	0.7
C	117	Oct-01	111	3	38	34	9.2	-19.0	7.7	0.4	-17.5	0.8

Mytilus californianus - A



Mytilus californianus - B






Figure 3. $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of *Mytilus californianus* periostracum samples, for individual animals A to C, in relation to the lip of the shell, commencing with the mantle edge at position 0. GCK positions are locations of growth checks interpreted to be annuli. Actual calendar years can be estimated starting with 1997, the year of sampling, between position 0 and the first annulus tick mark.

All three mussels had downward trending (towards origin) isotope values just prior to sampling – position 0 and adjacent samples. The $\delta^{13}\text{C}$ values for more recent years was above -18 .



The last three samples of each mussel had $\delta^{15}\text{N}$ less than $+8$. This was similar to the overall increase in ^{13}C content during the 1992 to 1997 period in the mussels sampled. There was no systematic shift in $\delta^{15}\text{N}$ across the years. There were, however, year-to-year shifts on the order of 1 ‰ that is similar to the isotopic gradient observed between PWS and the GOA and year-to-year differences observed in GOA stations (Kline 1999).

The fractional growth year plots, Figure 4, suggest large inter-annual differences in whether isotopic signatures increased, decreased or stayed the same. $\delta^{15}\text{N}$ values seem to start each growth period with values near $+8$ then diverge from that while $\delta^{13}\text{C}$ appear start with a more divergent range. Early divergent is consistent with the observed variations of GOA carbon during SEA since it was in May 1996 that $\delta^{13}\text{C}$ were very high. May is probably near the beginning of the growth period for *Mytilus* as growth rate is likely to be strongly influenced by seasonal warming such that spring growth is more likely to be slower per unit time than summer growth.



Further interpretation of inter-annual variation will require more sampling. This would be consistent with the bowhead whale baleen studies - over a decade of research was incorporated into Figure 1. Nonetheless, the inter-annual and seasonal isotopic variability suggest several levels of complexity in the ecological processes occurring in the Gulf.

Recent observations as part of on-going GLOBEC research appear to corroborate year-to-year inconsistencies in seasonal isotopic patterns observed in mussel periostracum (Kline unpublished). During 2001, $\delta^{15}\text{N}$ started low in March and April then increased in July and August whereas in 2002 it was uniformly high from May to October. During 2001, $\delta^{13}\text{C}$ started high in March, decreased to very low values in April then increased in July and August whereas in 2002 it gradually from a high in May to low values in October. The results thus far suggest greater levels of inter-annual variability than previous observed. Further research into the causes of isotopic shifts may reveal how ecosystem shifts take place.

Mytilus californianus - A



Mytilus californianus – B



Mytilus californianus – C




Figure 4. Fractional growth-year isotopic patterns. $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of *Mytilus californianus* periostracum samples, for individual animals A to C, in relation to apparent annual growth patterns. Values near 0 reflect early growth while values near 1, reflect growth just prior to an annulus. 1,2 ,and 3 of *Mytilus* ‘A’ corresponded to 1990, 1989, and 1988; 1,2 , 3, and 4 of *Mytilus* ‘B’ corresponded to 1989, 1987, and 1986, while 1, 2, 3, 4, and 5 of *Mytilus* ‘C’ corresponded to 1985, 1984, 1983, 1982, and 1981.

Literature cited

Anderson, P.J., J.E. Blackburn, and W.R. Bechtol. 1996. APEX Project Component L - Synthesis and analysis of Gulf of Alaska small-mesh trawl data 1953 to 1995. Abstracts of 1996 Restoration Projects. *Exxon Valdez Oil Spill Trustee Council*. Anchorage, Alaska.

Bidigare, R.R. and 15 other authors. 1997 Consistent fractionation of ^{13}C in nature and in the laboratory: Growth-rate effects in some haptophyte algae. *Global Biogeochemical Cycles*. 11:267-278.

Brodeur, R.D. and D.M. Ware. 1992. Long-term variability in zooplankton biomass in the subarctic Pacific Ocean. *Fish. Oceanogr.* 1:32-38.

Cooney, R.T. 1988. Distribution and ecology of zooplankton in the Gulf of Alaska: a synopsis. *In*: (Nemoto, T. and W.G. Pearcy (eds.) *The biology of the subarctic Pacific*, proceedings of the Japan-United States of America seminar on the biology of micronekton of the subarctic Pacific, Part I. *Bull. Ocean Res. Inst., University of Tokyo*. p. 27-41.

Cooney, R.T. 1993. A theoretical evaluation of the carrying capacity of Prince William Sound, Alaska, for juvenile Pacific salmon. *Fish. Res.* 18:77-87.

Francis, R.C. and S.R. Hare. 1994. Decadal-scale regime shifts in the large marine ecosystems of the northeast Pacific: a case for historical science. *Fish. Oceanogr.* 3:279-291.

Fry, B. 1988. Food web structure on Georges Bank from stable C, N, and S isotopic compositions. *Limnol. Oceanogr.* 33:1182-1190.

Laws, E. A., B. N. Popp, R. R. Bidigare, M. C. Kennicut, and S. A. Macko. 1995. Dependence of phytoplankton carbon isotopic composition on growth rate and $[\text{CO}_2]_{\text{aq}}$: Theoretical considerations and experimental results. *Geochem. Cosmochim. Acta*. 59:1131-1138.

Kline, Thomas Clayton, Jr. 1983. The effect of population density on the growth rate of the butter clam, *Saxidomus giganteus*. M. S. Thesis, University of Washington, Seattle, 104pp.

Kline, T. 1997a. Sound Ecosystem Assessment: Confirming food web dependencies in the Prince William Sound Ecosystem using stable isotope tracers, *Exxon Valdez Oil Spill Restoration Project Annual Report (Restoration Project 963201)*, Prince William Sound Science Center, Cordova, Alaska.

Kline, T.C. 1997b. Confirming forage fish food web dependencies in Prince William Sound using natural stable isotope tracers. *In: Forage Fishes in marine Ecosystems, Proceedings of the International Symposium on the Role of Forage Fishes in Marine Ecosystems*. Alaska Sea Grant College Program report No. 97-01. University of Alaska Fairbanks. p. 257-269.

Kline, T.C. 1998a. Carbon and Nitrogen Isotopic Composition of Prince William Sound Pelagic Biota Shift on Annual Time Scales: A Tool for Monitoring Changes in Oceanographic Forcing. *EOS Trans. AGU, 79(1), Ocean Sciences Meet. Suppl., OS53.*

Kline, T. C., Jr. 1999a. Monitoring changes in oceanographic forcing using the carbon and nitrogen isotopic composition of Prince William Sound pelagic biota. *In: Ecosystem Approaches for Fisheries Management*. University of Alaska Sea Grant, AK-SG-99-01, Fairbanks. p. 87-95.

Kline, T. C., Jr. 1999b. Temporal and Spatial Variability of $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ in pelagic biota of Prince William Sound, Alaska. *Can. J. Fish. Aquat. Sci.* 56 (Suppl. 1) 94-117.

Kline, T.C., Jr. 2001. Evidence of biophysical coupling from shifts in natural stable carbon and nitrogen isotopes in Prince William Sound, Alaska. *In: Kruse, G.H., N. Bez, A. Booth, M.W. Dorn, S. Hills, R.N. Lipcius, D. Pelletier, C. Roy, S.J. Smith, and D. Witherell (eds.), Spatial Processes and Management of Marine Populations*. University of Alaska Sea Grant, AK-SG-01-02, Fairbanks. p. 363-375.

Kline, T.C., Jr. 2001. Trophic position of Pacific herring in Prince William Sound, Alaska, based on their stable isotope abundance. *In: F. Funk, J. Blackburn, D. Hay, A.J. Paul, R. Stephenson, R. Toresen, and D. Witherell (eds.), Herring: Expectations for a New Millennium*. University of Alaska Sea Grant, AK-SG-01-04, Fairbanks. p. 69-80.

Kline, T. C. Jr. 2002. Relative trophic Position of *Cancer magister* megalopae. *In: (eds.) Crabs in Cold Water Regions: Biology, Management, and Economics*. University of Alaska Sea Grant, AK-SG-02-01, Fairbanks. p.645-649.

Kline, T. C., Jr. and Daniel Pauly. 1998. Cross-validation of trophic level estimates from a mass-balance model of Prince William Sound using $^{15}\text{N}/^{14}\text{N}$ data. *In: Funk, F., T.J. Quinn II, J. Heifetz, J.N. Ianelli, J.E. Powers, J.F. Schweigert, P.J. Sullivan, and C.-I. Zhang. (eds.) Fishery Stock assessment Models*. Alaska Sea Grant College Program Report No. AK-SG-98-01. University of Alaska Fairbanks. p. 693-702.

Kline, T. C. Jr., J. J. Goering, O. A. Mathisen, P. H. Poe and P. L. Parker. 1990. Recycling of elements transported upstream by runs of Pacific salmon: I. $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ evidence in Sashin Creek, southeastern Alaska. *Can. J. Fish. Aquat. Sci.* 47:136-144.

Kline, T.C. Jr., J.J. Goering, O.A. Mathisen, P.H. Poe, P.L. Parker, and R.S. Scalan. 1993. Recycling of elements transported upstream by runs of Pacific salmon: II. $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ evidence in the Kvichak River watershed, southwestern Alaska. *Can. J. Fish. Aquat. Sci.* 50:2350-2365.

Kline, T. C. Jr., W. J. Wilson, and J.J. Goering. 1998. Natural isotope indicators of fish migration at Prudhoe Bay, Alaska. *Can. J. Fish. Aquat. Sci.* 55:1494-1502.

Minagawa, M., and E. Wada. 1984. Stepwise enrichment of ^{15}N along food chains: Further evidence and the relation between $\delta^{15}\text{N}$ and animal age. *Geochim. Cosmochim. Acta* 48:1135-1140.

Oakey, T.A. and D. Pauly (eds). 1998. Trophic Mass-balance model of Alaska's Prince William Sound ecosystem, for the post-spill period 1994-1996. *Fisheries Centre Research Reports* 6(4). 144p.

Schell, D.M. 2000. Declining carrying capacity in the Bering Sea: Isotopic evidence from whale baleen. *Limnol. Oceanogr.* 45:459-462.

Vander Zanden, M.J., G. Cabana, and J.B. Rasmussen. 1997. Comparing trophic position of freshwater fish calculated using stable nitrogen isotope ratios ($\delta^{15}\text{N}$) and literature dietary data. *Can. J. Fish. Aquat. Sci.* 54:1142-1158.